BASELINE → TREATMENT → FOLLOW-UP

Treatment checks across the course

Relative stages, not equally spaced dates. Agree who books each check, reviews results and gives advice if a check is missed.

Check	Before starting	First treatment	During treatment	Phase / dose review	After treatment
Bloods and symptoms	Baseline blood count, kidney and liver tests; DPD before fluorouracil unless already tested	Confirm tests and prescription before the first dose	Blood count, kidney and liver tests before each cycle, with symptom review	Recheck after a hold or treatment change	Follow unresolved effects and the agreed disease-review plan
Treatment review	Confirm exact cancer setting and biomarkers	Explain expected benefit and how it will be assessed	Report new symptoms between visits	Review response, tolerability and active medicines	Agree the next treatment or surveillance plan
Electrolytes and EGFR effects	Check electrolyte and skin-risk plan	Skin care and prompt reaction reporting	Monitor magnesium, calcium, potassium, skin and eyes	Do not stop follow-up because the infusion stops	Monthly electrolytes, magnesium and calcium for two months after the last panitumumab dose
Heart assessment	Review cardiac history; baseline ECG if cardiac function is compromised	Report any new chest pain or breathing change	Chest pain, collapse or severe breathing difficulty needs emergency assessment	Additional heart tests only when clinically indicated	Follow any identified heart problem; no fixed routine echo series
INR only if taking warfarin	Tell the team about warfarin; arrange INR monitoring	INR as clinically indicated, rather than a universal weekly schedule	Report bleeding, bruising or significant changes in intake	The prescribing team adjusts warfarin if needed	Continue the individually agreed anticoagulation follow-up

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00448 v5 · official regimen

THE WHOLE TREATMENT JOURNEY

Panitumumab + FOLFIRI: treatment and checks

RAS wild-type metastatic colorectal cancer; first-line or selected second-line

A 14-day cycle: panitumumab, irinotecan, folinic acid and fluorouracil start on Day 1, followed by a fluorouracil pump for about 46 hours. Arrange disconnection around Day 3. Continue according to benefit and tolerance.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	First treatment	During treatment	Phase / dose review	After treatment
Bloods and symptoms NCCP 00448	Baseline blood count, kidney and liver tests; DPD before fluorouracil unless already tested	Confirm tests and prescription before the first dose	Blood count, kidney and liver tests before each cycle, with symptom review	Recheck after a hold or treatment change	Follow unresolved effects and the agreed disease-review plan
Treatment review NCCP 00448	Confirm exact cancer setting and biomarkers	Explain expected benefit and how it will be assessed	Report new symptoms between visits	Review response, tolerability and active medicines	Agree the next treatment or surveillance plan
Electrolytes and EGFR effects NCCP 00448	Check electrolyte and skin-risk plan	Skin care and prompt reaction reporting	Monitor magnesium, calcium, potassium, skin and eyes	Do not stop follow-up because the infusion stops	Monthly electrolytes, magnesium and calcium for two months after the last panitumumab dose
Heart assessment NCCP 00448	Review cardiac history; baseline ECG if cardiac function is compromised	Report any new chest pain or breathing change	Chest pain, collapse or severe breathing difficulty needs emergency assessment	Additional heart tests only when clinically indicated	Follow any identified heart problem; no fixed routine echo series
INR only if taking warfarin NCCP 00448	Tell the team about warfarin; arrange INR monitoring	INR as clinically indicated, rather than a universal weekly schedule	Report bleeding, bruising or significant changes in intake	The prescribing team adjusts warfarin if needed	Continue the individually agreed anticoagulation follow-up

What each check is for

1

A repeating 14-day cycle

Day ward → home pump → review

This combination aims to control bowel cancer that has spread and has a confirmed RAS wild-type result. The choice depends on previous treatment and your health.

Ask / act: Confirm the phase, prescription, tests and supportive medicines before starting.

Sources: NCCP 00448

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-04 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.33