

Modified FOLFOX-6

Gastrointestinal · NCCP 00209 · version 10a

Your medicines

Oxaliplatin · Folinic acid · 5-fluorouracil (5-FU)

Your treatment schedule

Every 14 days · See page 2 for the treatment phases.

A picture of side effects

Tell your team about new or worsening symptoms. The same symptom can have more than one cause. Numbers connect each symptom box to its location.

Oxaliplatin · nerve and cold effects

5-FU / chemotherapy · other effects

1 Mouth / throat

5-FU / oxaliplatin

Mouth ulcers or soreness; sudden throat tightness needs prompt assessment, including after cold exposure.

2 Heart / breathing

5-fluorouracil (5-FU)

New chest pain, pressure, fainting or breathlessness needs urgent help, including during the infusion.

3 Blood / infection

Oxaliplatin / 5-FU

Fever, shivers or sudden illness: contact oncology immediately.

4 Stomach / bowel

5-FU / oxaliplatin

Persistent diarrhoea, vomiting or inability to drink needs advice.

5 Hands / nerves

Oxaliplatin

Cold-triggered tingling or numbness; report symptoms that persist between cycles.

6 Feet / nerves

Oxaliplatin

Numbness, altered sensation or difficulty walking should be reported before the next dose.

7 Skin / mouth


5-fluorouracil (5-FU)

Report significant skin changes or mouth soreness that limits eating or drinking.

8 Line / home pump

46-hour 5-FU infusion

Call for leakage, disconnection, pain, redness, swelling or a pump alarm. Follow local pump instructions.



This map shows selected effects to discuss; it does not show their likelihood or every possible side effect. Blood-count and whole-body effects are marked schematically.

Selected symptom patterns for discussion. Tell your oncology team about any new or worsening symptom. See page 2 for medicine-specific effects, monitoring and the portal QR.

Modified FOLFOX-6

Gastrointestinal · NCCP 00209 · version 10a

TREATMENT JOURNEY

Day 1 Oxaliplatin and folinic acid are given by drip, then 5-FU is started.	Days 1-3 A portable pump gives 5-FU continuously for about 46 hours.	Around day 3 Pump disconnect. Follow your local line and pump instructions.	Days 4-14 Recovery and monitoring. Symptoms can occur on any day.	Day 15 Next cycle if review and blood tests allow.
---	--	---	---	--

IMPORTANT MEDICINE-SPECIFIC EFFECTS

Oxaliplatin

- Cold-triggered tingling or throat/jaw sensations can occur soon after treatment.
- Persistent peripheral neuropathy can develop with cumulative treatment.
- Infusion hypersensitivity can occasionally be severe.

5-fluorouracil (5-FU)

- Severe toxicity can occur with reduced DPD activity.
- Diarrhoea and mouth/gut inflammation can become severe.
- New chest pain or rhythm symptoms require urgent assessment; hand-foot soreness can occur.

Other / shared treatment effects: The anatomy page highlights important symptom patterns for this exact regimen, but it cannot show every possible effect. Fatigue, nausea, appetite change, mouth/bowel symptoms or blood-count changes may occur depending on the medicines used. Report new or worsening symptoms even if they are not listed here.

Monitoring / medicines

- DPD status is checked before fluoropyrimidine treatment.
- Before each cycle: blood count, kidney/liver bloods and assessment of nerve symptoms.
- Follow your own labelled anti-sickness, mouth-care and diarrhoea medicines.

Bring / track

- Your current medicines, including non-prescription products.
- A note of new symptoms, when they started and what they stopped you doing.
- Any temperature, blood-pressure or bowel-count readings your team asked you to record.
- Questions about the purpose of treatment, scan/review timing or changes to the plan.

WHEN TO CALL - DO NOT WAIT FOR THE NEXT APPOINTMENT

Call your oncology SOS service for fever or shivers, sudden illness, 4 or more diarrhoea episodes in a day, vomiting despite medicines, mouth sores stopping fluids, bleeding, new chest pain or breathlessness. Severe chest pain or breathing difficulty: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

Consent discussion support

Use the approved consent process to document treatment aim, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

Regimen information only; this QR contains no patient details.

<https://sactcheck.com/patient/00209/>

Official NCCP 00209 v10a · SACTCheck v0.81.2 · patient information prototype for clinical review.

