

TCHP

Breast · NCCP 00722 · version 2

Your medicines

Docetaxel · Carboplatin · Trastuzumab · Pertuzumab

Your treatment schedule


Every 21 days · See page 2 for the treatment phases.

A picture of side effects

Tell your team about new or worsening symptoms. The same symptom can have more than one cause. Numbers connect each symptom box to its location.

Trastuzumab / pertuzumab · HER2 treatment

Docetaxel / carboplatin · chemotherapy



This map shows selected effects to discuss; it does not show their likelihood or every possible side effect. Blood-count and whole-body effects are marked schematically.

Selected symptom patterns for discussion. Tell your oncology team about any new or worsening symptom. See page 2 for medicine-specific effects, monitoring and the portal QR.

TCHP

Breast · NCCP 00722 · version 2

TREATMENT JOURNEY

Before cycle 1 Baseline blood tests and cardiac assessment are reviewed.	Cycle day 1 Docetaxel + carboplatin + trastuzumab + pertuzumab are given.	Days 2-21 Recovery, supportive medicines and symptom monitoring.	Every 21 days The cycle repeats if review and blood tests allow.	After chemotherapy HER2-directed treatment usually continues according to the post-operative/adjuvant treatment plan.
--	---	--	--	---

IMPORTANT MEDICINE-SPECIFIC EFFECTS

Docetaxel - Severe neutropenia/infection can occur. - Fluid retention, swelling, nerve symptoms, nail/skin changes and infusion reactions can occur.	Carboplatin - Low blood counts, especially neutropenia and low platelets, can occur. - Allergic infusion reactions can occur, particularly after repeated exposure. - Nausea and vomiting may occur despite preventative medicines.
Trastuzumab + pertuzumab - Heart function can fall and is monitored during HER2-directed treatment. - Infusion/administration reactions can occur. - Pertuzumab commonly causes diarrhoea; rare serious lung reactions can occur with HER2 therapy.	

Other / shared treatment effects: The anatomy page highlights important symptom patterns for this exact regimen, but it cannot show every possible effect. Fatigue, nausea, appetite change, mouth/bowel symptoms or blood-count changes may occur depending on the medicines used. Report new or worsening symptoms even if they are not listed here.

Monitoring / medicines - Follow the prescribed steroid premedication around docetaxel. - Growth-factor support may be part of the plan; follow the exact product and timing given by your oncology team. - Cardiac function is monitored because trastuzumab/pertuzumab can affect the heart.	Bring / track - Your current medicines, including non-prescription products. - A note of new symptoms, when they started and what they stopped you doing. - Any temperature, blood-pressure or bowel-count readings your team asked you to record. - Questions about the purpose of treatment, scan/review timing or changes to the plan.
--	--

WHEN TO CALL - DO NOT WAIT FOR THE NEXT APPOINTMENT
Call oncology urgently for fever, severe diarrhoea, new breathlessness, swelling, severe rash, uncontrolled vomiting, significant bleeding or a rapidly worsening symptom. Severe breathing difficulty or chest pain: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

Consent discussion support

Use the approved consent process to document treatment aim, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

Regimen information only; this QR contains no patient details.

<https://sactcheck.com/patient/00722/>

Official NCCP 00722 · SACTCheck v0.81.2 · patient information prototype for clinical review.

