BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Columns show sequence, not equal lengths. Count 12 weeks from first HER2 treatment (after AC where applicable): around weekly antibody dose 13 or q21-day cycle 5. Preserve elapsed time across later phase / route changes and delays. The treating team owns booking, result review and action when a test is overdue.

Check	Before starting	Chemotherapy visits	Phase change / 12 weeks	Continuation / local care	Completion / follow-up
Blood tests	Baseline FBC, renal and liver profile	FBC, renal and liver profile before each cycle. Weekly-day checks follow the active regimen and local delivery policy.	Reassess before a new phase	Companion drugs / abnormal results determine additional tests	Follow persistent abnormalities and the next plan
Echo / cardiac assessment	Baseline before HER2 treatment.	Earlier assessment for symptoms or cardiac concern	Count 12 weeks from first HER2 treatment (after AC where applicable): around weekly antibody dose 13 or q21-day cycle 5. Preserve elapsed time across later phase / route changes and delays.	Every 12 weeks during HER2 treatment, including continuation.	After year 1, six-monthly echo may be considered only for selected asymptomatic low/ moderate-risk metastatic patients with normal assessments (ESC). Confirm a team-approved plan.
ECG / rhythm	Baseline ECG as part of trastuzumab cardiac assessment	Symptoms, interacting medicines or clinical indication	An ECG does not replace an echo	Repeat if symptoms or the cardiac plan require it	Agree follow-up if abnormal
Imaging / markers	Document baseline disease and purpose of scans	Review symptoms and response	Metastatic: scan if due	For metastatic breast disease, response assessment commonly every 2–4 months, adapted to sites, symptoms and treatment. Markers can supplement assessment; they do not replace imaging or alone determine a treatment change.	At a treatment change, document disease status and the next imaging plan. Tumour markers supplement imaging.
Bone health / DEXA	Assess endocrine treatment and fracture risk	Not a routine scan at each cycle	Baseline DEXA when indicated	DEXA before aromatase inhibitor / ovarian suppression when indicated; repeat according to fracture risk, results and endocrine plan.	Risk- and result-based follow-up

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00204 v11 · official source · ESMO 2024 early breast cancer guideline · treatment and follow-up · ESC 2022 cardio-oncology guideline · cardiac surveillance and risk · NICE NG101 · bone health and follow-up · ESMO metastatic breast cancer guideline · 2026 · EMA Herceptin product information

THE WHOLE TREATMENT JOURNEY

Docetaxel + pertuzumab + trastuzumab: the whole pathway

HER2-positive metastatic or unresectable locally recurrent breast cancer

IV pertuzumab, IV trastuzumab and docetaxel every 21 days. Docetaxel duration is reviewed separately; the two HER2 antibodies may continue after chemotherapy stops while beneficial and tolerated.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before systemic treatment	Chemotherapy phase	Next phase / reassessment	Local / continuing treatment	Follow-up / change
Cancer medicines NCCP 00204	Confirm indication, organ function and access	Taxane ± HER2 treatment	Review tolerance and response	HER2 continuation under the agreed plan	Review outcome and next treatment
Surgery METASTATIC	Site-specific procedures if indicated	No automatic surgery during these cycles	Reassess local-treatment needs	Individual multidisciplinary plan	Local follow-up
Radiotherapy METASTATIC	Assess indication	Only when clinically indicated	Coordinate surgery and recovery	Radiation team agrees timing	Assess late effects / local follow-up
Blood tests NCCP 00204	Baseline FBC, renal, liver	FBC, renal and liver profile before each cycle. Weekly-day checks follow the active regimen and local delivery policy.	Review before transition	Continue according to active medicines	Follow abnormalities
Heart tests NCCP 00204 · SMPC · ESC	Baseline echo / MUGA	Symptoms can prompt earlier tests	At 12 weeks from HER2 start	Every 12 weeks / 3 months during HER2	Agree post-treatment cardiac plan
Scans / markers EARLY · METASTATIC	Baseline staging as indicated	Clinical response / symptoms	Response scan if due	Response imaging if metastatic	Symptoms can trigger earlier tests
Endocrine / bone health EARLY · BONE	Check receptors / endocrine indication	Discuss the later plan	Start / continue when agreed	May overlap HER2 / RT if appropriate	Longer-term bone health and endocrine review

What each check is for

1 Blood checks

FBC, renal and liver profile before each cycle. Weekly-day checks follow the active regimen and local delivery policy.

Results help the team judge treatment safety.

Ask / act: Confirm the next test and who reviews it.

Sources: NCCP 00204

2 Heart assessment

Baseline before HER2 treatment; repeat every 12 weeks during treatment. Repeat at completion; 4–8-week checks may be needed with impairment.

Some heart changes occur without symptoms.

Ask / act: Keep the appointment and report new symptoms between scans.

Sources: NCCP 00204 · SMPC · ESC

3 Imaging and markers

For metastatic breast disease, response assessment commonly every 2–4 months, adapted to sites, symptoms and treatment. Markers can supplement assessment; they do not replace imaging or alone determine a treatment change.

Each scan should answer a defined clinical question.

Ask / act: Ask when the result will be reviewed.

Sources: EARLY · METASTATIC

4 Booking and handover

At initiation, phase changes and completion

Expected tests can be missed without an owner.

Ask / act: Record booking owner, result reviewer, due window and escalation contact.

Sources: NCCP 00204

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-10-17 · Source reconciled; independent clinical, pharmacy and patient review pending · SACTCheck v0.84.0