AC · every 3 weeks

NCCP 00252 · Source reconciled; independent clinical, pharmacy and patient review pending

PRACTICAL TREATMENT GUIDE

Your practical treatment plan

Doxorubicin and cyclophosphamide on Day 1 every 21 days for four cycles. Any subsequent taxane follows a separately agreed regimen.

STEP 1

Before the first visit

Blood tests, relevant cardiac assessment, medicines review, fertility discussion and the consent conversation come first.

STEP 2

Plan venous access

A cannula, PICC or implanted port may be used. If a port is chosen, a trained team inserts it in a separate procedure with local anaesthetic and sometimes sedation; the team explains fasting, blood-thinner and wound instructions.

STEP 3

Infusion day

Staff check the line, give prescribed premedicines and treatment, and observe for reactions. Tell them immediately about burning, pain, rash or breathlessness.

STEP 4

Home and next review

This regimen does not routinely need a take-home infusion pump. Staff disconnect the infusion; a port is de-accessed when appropriate. Keep line-care and next-visit instructions.

Port insertion and removal

The line team explains placement, dressing care and when normal activities can resume. A port is accessed with a special needle during treatment. Removal of the implanted device is a separate planned procedure when no longer needed; do not remove or adjust it yourself.

PICC care and urgent line symptoms

A PICC usually needs regular dressing and flushing appointments even between treatments. Fever, redness, swelling, pain, leakage or damage requires prompt advice. Your hospital supplies the exact care instructions.

Visit time and transport

First visits can take longer because of loading doses, premedicines and observation. Ask the unit about expected time, transport and any escort needed after sedation or line insertion.

Phase-specific visit pattern

Infusion and observation times differ by drug and first versus later doses; the unit confirms the full visit duration, including blood review and preparation.

Sources: NCCP 00252 v7 · official source · eviQ · implanted port patient information · eviQ · PICC patient information. Source reconciled; independent clinical, pharmacy and patient review pending.