# Docetaxel 100 mg/m² · every 3 weeks

Docetaxel 100 mg/m² by IV infusion on Day 1 every 21 days, with dose changes when needed. The number of cycles follows benefit, tolerance and the agreed treatment plan.

Source reconciled; independent clinical, pharmacy and patient review pending. Selected effects; not a complete medicine leaflet.

## A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

**1 Hands / feet**

Docetaxel

Report numbness or tingling before the next dose.

**3 Hair / scalp**


Chemotherapy phase

Hair loss is common. Ask about scalp cooling before treatment where suitable.

**5 Mouth**

Chemotherapy phase

Painful ulcers or difficulty drinking need advice.




**2 Blood / infection**

Chemotherapy phase

Fever, shivers or suddenly unwell: call immediately.

**4 Fluid / skin / nails**

Docetaxel

Report swelling, painful skin or nail changes.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects; the full risk discussion includes hair loss, fertility and other individual risks.

Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

# Your treatment, at home.

## Important medicine-specific effects

### Docetaxel

- Low blood counts, infection and sepsis
- Hair loss, mouth soreness and fatigue
- Numbness / tingling; sometimes lasting
- Fluid retention and nail changes
- Allergic infusion reaction

**Surgery, radiation and monitoring:** open the whole-treatment map or scan the QR.

## Make the plan practical

### The correct phase prescription

Taxanes use low-risk antiemetic support; hypersensitivity premedication is a separate requirement.

### Premedicines and growth factors

Docetaxel dexamethasone is taken for three days starting the day before treatment. Ask the team to reconcile this with antiemetic steroids.

### Tests and appointments

The team should name the booking owner, due window and person reviewing each required test. Tell them if an expected appointment has not arrived.

### After chemotherapy

Discuss recovery, fertility, rehabilitation and the next treatment or surveillance step.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

CUH · MON–FRI 8AM–5PM

**087 181 9645**

Acute Oncology Service

CUH · WEEKENDS / BANK HOLIDAYS 9AM–5PM

**021 4922000**

Ask for medical oncology

EVENINGS / OVERNIGHT

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

**Complete reference, visual timeline and passport**

https://sactcheck.com/patient/00202/

Sources and review status are available online. Follow your own prescription and hospital instructions.