NCCP 00260 · clinician draft

AC every 3 weeks → weekly paclitaxel

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Taxane / combination phase

Active medicines: Paclitaxel

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally	On treatment day Review QT, constipation and hepatic dosing; reconcile any steroid already prescribed
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days: duration to be confirmed Assess contraindications, renal function and interactions

Additional phase-specific support

- First paclitaxel dose: dexamethasone 8 mg IV, chlorphenamine 10 mg IV and famotidine 20 mg IV, each 30 minutes before treatment (NCCP suggested table).
- From Day 8 onward: chlorphenamine 10 mg IV and famotidine 20 mg IV 30 minutes before treatment. Routine dexamethasone may be omitted after the first dose; prior reaction or increased risk requires consultant review.
- Famotidine may be omitted under the consultant/local plan if appropriate. A prior reaction requires a separate rechallenge assessment; do not reuse the standard draft automatically.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00260 v7 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-17 · SACTCheck v0.84.0 · Regimen-level drafts; prescriber and oncology-pharmacy verification required