SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

Trastuzumab SC · early breast cancer

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

HER2-positive early breast cancer · SC component · NCCP 00285 · version 4 · 21-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
Trastuzumab under the skin is part of treatment to reduce recurrence in HER2-positive early breast cancer. The wider plan can include chemotherapy, surgery, radiotherapy and endocrine treatment when indicated.

Schedule
Trastuzumab 600 mg under the skin every 21 days, without a loading dose, to complete one year total. Previous IV or combination-treatment trastuzumab counts towards that year; a route switch does not start a new year.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Duration and treatment setting discussed
- ☐ Supportive medicines and urgent contact advice discussed
- ☐ Common, serious and lasting agent-specific risks discussed below
- ☐ Correct HER2 medicine and formulation confirmed
- ☐ Heart and other required monitoring explained
- ☐ Pregnancy / contraception / breastfeeding discussed where relevant
- ☐ Companion cancer medicines reviewed using their own information

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Trastuzumab
HER2-TARGETED ANTIBODY

- ☐ Heart pumping-function reduction / heart failure
- ☐ Allergic or administration-related reactions
- ☐ Injection-site pain, redness or swelling
- ☐ Diarrhoea, fatigue and rare serious lung effects

Discuss expected benefit, alternatives, route, duration, cardiac monitoring, pregnancy / contraception, urgent contacts and individual treatment risks. Add any companion drug using its own regimen information.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00285 v4.

SACTCheck v0.83.9
Developed by Dr Paul O'Brien