NCCP 00605 · VERSION 2

Talazoparib · breast monotherapy

Talazoparib is taken once daily continuously until disease progression or unacceptable toxicity. The usual starting dose is 1 mg; kidney function, interactions and side effects can require a different prescription.

Oral PARP inhibitor · selected medicine-specific effects. Full details and sources are in the complete reference.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Blood / infection


Talazoparib

Fever, infection or unusual bruising needs prompt advice.

2 Energy / breathing

Talazoparib

Marked fatigue or breathlessness may reflect anaemia.



3 Nausea / appetite

Talazoparib

Call if vomiting or poor intake prevents fluids or medicines.

4 Hair / scalp

Talazoparib

Hair thinning or loss can occur; ask about practical support.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact oncology urgently for fever, shivers, infection, unusual bleeding, marked fatigue or new breathlessness. Severe chest pain, major breathing difficulty, collapse or stroke symptoms: 112 / 999.

YOUR PRACTICAL TREATMENT PLAN

Know the next step.

Taking treatment

Talazoparib is taken once daily continuously until disease progression or unacceptable toxicity. The usual starting dose is 1 mg; kidney function, interactions and side effects can require a different prescription.

Swallow capsules whole with water, with or without food. Do not open or dissolve them. If a dose is missed or vomited, take the next prescribed dose at its usual time; do not take a replacement or double dose.

Checks and follow-up

Check FBC, kidney and liver function before starting and every four weeks. More frequent blood review may be needed for low counts; prolonged abnormalities require specialist assessment. Confirm the germline BRCA result and previous treatment. Review pregnancy status and contraception, kidney function and interacting medicines, particularly P-glycoprotein inhibitors.

The whole-treatment map shows the checks for this exact treatment setting.

Support between reviews

Blood counts

Keep four-weekly blood reviews and any additional tests. Fever or unusual bleeding needs prompt advice.

Taking capsules

Use the strength and number on your current prescription. Do not replace a missed or vomited dose.

Nausea

Use the prescribed rescue plan; call if vomiting affects fluids or tablets.

Medicines review

Check every new medicine or supplement with the oncology pharmacist.

Oncology daytime: Insert numbers here. **Out of hours:** Insert numbers here. Follow your own hospital’s urgent-care plan. Emergency: 112 / 999.

Your complete treatment reference

https://sactcheck.com/patient/00605/

Public information only. No personal information is encoded.