BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Relative milestones, not equal time intervals. Your team books tests and reviews results; this map does not confirm completion.

Check	Before starting	First treatment	During treatment	Change / completion	Follow-up
Blood / organ function	FBC, renal, liver baseline	Review results before dose	FBC / renal / liver every cycle	Confirm counts for each planned dose	Investigate persisting problems
Heart / nerves	Cardiac function if history / prior anthracycline	Neuropathy and symptom review	Cardiac review when clinically indicated	Assess response after two cycles	Report persistent symptoms

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00230 v7 · official regimen · NICE NG101 · bone health and follow-up · ASCO · biomarkers in metastatic breast cancer · ESMO metastatic breast cancer guideline · 2026

THE WHOLE TREATMENT JOURNEY

Nab-paclitaxel · every 3 weeks: treatment and checks

Metastatic breast cancer after first-line failure; standard anthracycline therapy unsuitable

Nab-paclitaxel on Day 1 every 21 days. Review response after two cycles; the NCCP protocol says to discontinue if there is no response. Further treatment depends on benefit and tolerance.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	First treatment	During treatment	Change / completion	Follow-up
Treatment plan NCCP 00230	Baseline eligibility / tests	Day 1 infusion	Review response after two cycles	Repeat q21 days if benefiting / tolerating	Next treatment plan
Blood / organ function NCCP 00230	FBC, renal, liver baseline	Review results before dose	FBC / renal / liver every cycle	Confirm counts for each planned dose	Investigate persisting problems
Heart / nerves NCCP 00230	Cardiac function if history / prior anthracycline	Neuropathy and symptom review	Cardiac review when clinically indicated	Assess response after two cycles	Report persistent symptoms
Cancer review METASTATIC · MARKERS	Confirm setting	Agree review plan	Response / symptom review	Next treatment / completion	Agreed surveillance

What each check is for

1 Before treatment

Before the first dose

Metastatic breast cancer after first-line failure; standard anthracycline therapy unsuitable. FBC, kidney and liver tests before treatment and each cycle. Review response after two cycles; the source specifies stopping if there is no response.

Ask / act: Confirm eligibility, test results and medicine list.

Sources: NCCP 00230

2 Other checks

At the specified review points

Baseline cardiac assessment for a significant cardiac history or prior anthracycline exposure; further assessment as clinically indicated. Review neuropathy before doses.

Ask / act: Confirm who arranges and reviews each check.

Sources: NCCP 00230

3 Support and practical plan

At each treatment or phase change

Routine allergy premedicines are usually not required for this albumin-bound formulation. Low-risk antiemetic support is separate; previous reactions may change the plan.

Ask / act: Choose the matching support plan; confirm active medicines.

Sources: NCCP 00230

4 Response and follow-up

Agreed review points

Agree the scan type and review interval for the known disease sites. Review response, symptoms and toxicity together; new or worsening symptoms may need an earlier assessment. Keep the agreed follow-up and report persistent or new symptoms between visits.

Ask / act: Review symptoms, results and patient priorities.

Sources: METASTATIC · MARKERS

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-10-25 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.19