Nivolumab, ipilimumab, paclitaxel and carboplatin

NCCP 00712 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Visits and duration

Cycle 1 lasts 42 days: nivolumab, ipilimumab, paclitaxel and carboplatin on Day 1; nivolumab, paclitaxel and carboplatin on Day 22. These are the only two planned chemotherapy administrations. From Cycle 2, nivolumab is given on Days 1 and 22 and ipilimumab on Day 1. Continue while appropriate until progression, unacceptable toxicity or up to 24 months without progression. Your team reviews the plan and may delay or stop treatment.

STEP 2

Time in the unit

Nivolumab over 30 minutes, then ipilimumab over 30 minutes on Day 1 only. Paclitaxel over three hours, then carboplatin over 30 minutes on both Cycle 1 treatment days. Ipilimumab vital signs are checked every 30 minutes during infusion and for an hour afterwards. Blood tests, preparation and observation add to time in the unit.

STEP 3

Medicines and monitoring

Blood count, kidney and liver tests before each treatment; glucose and thyroid tests before each 42-day cycle. Cortisol and neuropathy checks when indicated. Paclitaxel premedication includes corticosteroid and antihistamine treatment, with H2-blocker decisions made by the team. The source allows consultant-directed changes; ask for a written plan for each of the two chemotherapy visits.

STEP 4

Transport and support

Arrange travel and help if needed. Keep urgent contacts and alert information with you.

Sources: NCCP 00712 v3a · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.