NCCP 00849 · clinician draft

Nivolumab, gemcitabine and carboplatin before surgery

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Day 8 · gemcitabine alone

Active medicines: gemcitabine

Emetogenic risk: low

Medicine	Dose / route	When
Dexamethasone	8 mg orally	Day 8 before gemcitabine if prescribed NCIS low-risk reference; individualise and reconcile with any other steroid course.

Additional phase-specific support

- Separate Day 8 blood-count and symptom review. Do not repeat Day 1 high-risk prophylaxis automatically.
- Nivolumab can inflame healthy organs during treatment or after it stops. Report new symptoms promptly and carry your immune alert card for at least five months after the last dose.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00849 v2 · pNCCP 00310; NCIS V8 Table 1 · NCCP NCIS antiemetics V8

Sources checked 2026-10-29 · SACTCheck v0.84.27 · Regimen-level drafts; prescriber and oncology-pharmacy verification required