NCCP 00712 · clinician draft

Nivolumab, ipilimumab, paclitaxel and carboplatin

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Cycle 2 onwards · Day 1 nivolumab + ipilimumab

Active medicines: nivolumab + ipilimumab

Emetogenic risk: low

Medicine	Dose / route	When
Ondansetron	8 mg orally	Day 1 before treatment, if following the NCIS low-risk default Individualise/local policy. No automatic antiemetic corticosteroid course.
Metoclopramide	10 mg orally three times daily as needed	Subsequent days as prescribed Specify duration and check contraindications/interactions.

Additional phase-specific support

- Both immune medicines can inflame healthy organs during treatment or after it stops. Keep your immune alert card and report new symptoms promptly, including for at least five months after the last dose.
- Chemotherapy is complete; review and stop induction prophylaxis rather than automatically repeating it.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00712 v3a · pNCCP 00712 pp8–9; NCIS V8 Table 1 p4 · NCIS low-risk row

Sources checked 2026-10-29 · SACTCheck v0.84.27 · Regimen-level drafts; prescriber and oncology-pharmacy verification required