NCCP 00714 · clinician draft

Nivolumab, ipilimumab, pemetrexed and cisplatin

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Cycle 2 onwards Day 22 · nivolumab alone, between ipilimumab visits

Active medicines: nivolumab

Emetogenic risk: Minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- No routine prophylactic antiemetic course for nivolumab-only minimal-risk treatment; prescribe symptom-directed rescue under local policy if needed.
- Ipilimumab is due at the next 42-day cycle Day 1, not every nivolumab visit.
- Immune toxicity can still occur between ipilimumab visits; the joint hold rule remains relevant.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00714 v4 · pNCCP 00714 schedule and emetogenic potential · Current NCCP regimen

Sources checked 2026-10-29 · SACTCheck v0.84.27 · Regimen-level drafts; prescriber and oncology-pharmacy verification required