Nivolumab and ipilimumab for pleural mesothelioma

NCCP 00792 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Visits and duration

Each cycle lasts 42 days: nivolumab and ipilimumab on Day 1, then nivolumab alone on Day 22. Nivolumab is every three weeks and ipilimumab every six weeks. This regimen contains no chemotherapy. Continue while appropriate until progression, unacceptable toxicity or up to 24 months without progression. Your team reviews the plan and may delay or stop treatment.

STEP 2

Time in the unit

Nivolumab over 30 minutes, then ipilimumab over 30 minutes on Day 1 only. Ipilimumab vital signs are checked every 30 minutes during infusion and for an hour afterwards. Blood tests, preparation and observation add to time in the unit.

STEP 3

Medicines and monitoring

Blood count, kidney, liver, glucose and thyroid tests before each 42-day cycle; symptoms reviewed before each infusion and additional tests when indicated. Cortisol if clinically indicated. Carry treatment details and the immune alert card. Describe your usual bowel pattern, breathing, energy, skin and hormone symptoms, and tell the team about autoimmune illness, infections and all medicines.

STEP 4

Transport and support

Arrange travel and help if needed. Keep urgent contacts and alert information with you.

Sources: NCCP 00792 v2 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.