NCCP 00792 · clinician draft

# Nivolumab and ipilimumab for pleural mesothelioma

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

## Every 42-day cycle · Day 1 nivolumab + ipilimumab

**Active medicines:** nivolumab + ipilimumab

**Emetogenic risk:** low

| Medicine | Dose / route | When |
|---|---|---|
| Ondansetron | 8 mg orally | Day 1 before treatment, if following the NCIS low-risk default<br>Individualise/local policy. No automatic antiemetic corticosteroid course. |
| Metoclopramide | 10 mg orally three times daily as needed | Subsequent days as prescribed<br>Specify duration and check contraindications/interactions. |

### Additional phase-specific support

- Both immune medicines can inflame healthy organs during treatment or after it stops. Keep your immune alert card and report new symptoms promptly, including for at least five months after the last dose.

**Before signing:** verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00792 v2 · pNCCP 00792 p5; NCIS V8 Table 1 p4 · NCIS low-risk row

Sources checked 2026-10-29 · SACTCheck v0.84.27 · Regimen-level drafts; prescriber and oncology-pharmacy verification required