SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Cetuximab + carboplatin + fluorouracil

Treatment of locally advanced or metastatic squamous cell cancer of the head and neck when cisplatin cannot be used or tolerated. · NCCP 00418 · version 7 · 21-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
Cetuximab with carboplatin and fluorouracil is used to treat squamous head and neck cancer when cisplatin is unsuitable. Your team explains the treatment aim, possible benefit and alternatives for your situation.

Schedule
Carboplatin is given on Day 1, fluorouracil on Days 1–4, and cetuximab on Days 1, 8 and 15, for up to 6 cycles every 21 days. Review determines what happens afterwards. Your team confirms any treatment after the combination phase; continuing cetuximab needs an agreed prescription and calendar.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ DPD result and severe fluoropyrimidine risks reviewed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Cetuximab
EGFR-TARGETED ANTIBODY

- ☐ Infusion reactions, including anaphylaxis
- ☐ Acne-like rash, dry or cracked skin and nail infections
- ☐ Low magnesium and other mineral changes
- ☐ Eye inflammation
- ☐ Diarrhoea and dehydration
- ☐ Rare serious lung inflammation

Carboplatin
PLATINUM CHEMOTHERAPY

- ☐ Low blood counts and serious infection
- ☐ Low platelets and bleeding
- ☐ Anaemia and fatigue
- ☐ Nausea and vomiting
- ☐ Infusion or allergic reactions
- ☐ Kidney, nerve or hearing effects

Fluorouracil
ANTIMETABOLITE CHEMOTHERAPY

- ☐ Low blood counts and serious infection
- ☐ Mouth and throat soreness
- ☐ Diarrhoea and dehydration
- ☐ Painful palms or soles
- ☐ Heart problems including chest pain
- ☐ Rare severe neurological or early excessive toxicity

Discuss the individual aim, expected benefit, alternatives and uncertainties; common, serious and lasting effects; infusion reactions and infection; fertility and pregnancy; treatment burden; support medicines and urgent contacts. For a locally advanced treatment plan, confirm whether cetuximab use falls outside the current licensed combination indication and discuss this explicitly.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00418 v7.

SACTCheck v0.84.47
Developed by Dr Paul O'Brien