SACTCHECK · TREATMENT PASSPORT

Carboplatin + fluorouracil after nasopharyngeal chemoradiotherapy

A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

After recovery, adjuvant chemotherapy is planned to begin 4 weeks after completion of weekly carboplatin with radiotherapy (National Cancer Control Programme (NCCP) 00419). Carboplatin is given on Day 1 and fluorouracil over Days 1–4, every 28 days for up to 3 cycles. The treating team confirms readiness and dates.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00552/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending

NCCP 00552 V6 · ONE CYCLE, ONE PAGE

Carboplatin + fluorouracil after nasopharyngeal chemoradiotherapy

Cycle number: _____ Start date: _____ Next review: _____

Record the date, medicines actually received, delays or omissions, test results and concerns with your team. Use a separate record for any changed treatment phase.

Adjuvant cycle 1 of up to 3 · copy this sheet for another cycle of the same phase.

Carboplatin · Platinum chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Fever / chills	☐ No fever or chills	☐ New chills / feeling unwell: call	☐ Fever: use the written urgent plan now	☐ Fever with marked weakness: urgent assessment	☐ Collapse / confusion / breathing difficulty: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999
Infusion symptoms	☐ None	☐ New flushing, chills or itching: tell the nurse immediately	☐ Symptoms during infusion: tell the nurse immediately	☐ Wheeze, swelling or faintness: immediate urgent help	☐ Severe breathing difficulty or collapse: emergency help
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help

Fluorouracil · Antimetabolite chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Heart symptoms	☐ None	☐ New palpitations: contact team promptly	☐ Palpitations with dizziness: urgent assessment	☐ Chest pain or fainting: emergency assessment	☐ Severe chest pain, collapse or breathing difficulty: 112/999
Fever / chills	☐ No fever or chills	☐ New chills / feeling unwell: call	☐ Fever: use the written urgent plan now	☐ Fever with marked weakness: urgent assessment	☐ Collapse / confusion / breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

NCCP 00552 V6 · ONE CYCLE, ONE PAGE

Carboplatin + fluorouracil after nasopharyngeal chemoradiotherapy

Cycle number: _____ Start date: _____ Next review: _____

Record the date, medicines actually received, delays or omissions, test results and concerns with your team. Use a separate record for any changed treatment phase.

Adjuvant cycle 2 of up to 3 · copy this sheet for another cycle of the same phase.

Carboplatin · Platinum chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Fever / chills	☐ No fever or chills	☐ New chills / feeling unwell: call	☐ Fever: use the written urgent plan now	☐ Fever with marked weakness: urgent assessment	☐ Collapse / confusion / breathing difficulty: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999
Infusion symptoms	☐ None	☐ New flushing, chills or itching: tell the nurse immediately	☐ Symptoms during infusion: tell the nurse immediately	☐ Wheeze, swelling or faintness: immediate urgent help	☐ Severe breathing difficulty or collapse: emergency help
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help

Fluorouracil · Antimetabolite chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Heart symptoms	☐ None	☐ New palpitations: contact team promptly	☐ Palpitations with dizziness: urgent assessment	☐ Chest pain or fainting: emergency assessment	☐ Severe chest pain, collapse or breathing difficulty: 112/999
Fever / chills	☐ No fever or chills	☐ New chills / feeling unwell: call	☐ Fever: use the written urgent plan now	☐ Fever with marked weakness: urgent assessment	☐ Collapse / confusion / breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

NCCP 00552 V6 · ONE CYCLE, ONE PAGE

Carboplatin + fluorouracil after nasopharyngeal chemoradiotherapy

Cycle number: _____ Start date: _____ Next review: _____

Record the date, medicines actually received, delays or omissions, test results and concerns with your team. Use a separate record for any changed treatment phase.

Adjuvant cycle 3 of up to 3 · copy this sheet for another cycle of the same phase.

Carboplatin · Platinum chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Fever / chills	☐ No fever or chills	☐ New chills / feeling unwell: call	☐ Fever: use the written urgent plan now	☐ Fever with marked weakness: urgent assessment	☐ Collapse / confusion / breathing difficulty: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999
Infusion symptoms	☐ None	☐ New flushing, chills or itching: tell the nurse immediately	☐ Symptoms during infusion: tell the nurse immediately	☐ Wheeze, swelling or faintness: immediate urgent help	☐ Severe breathing difficulty or collapse: emergency help
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help

Fluorouracil · Antimetabolite chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Hand-foot soreness	☐ None	☐ Redness or tingling without pain	☐ Pain limits usual tasks: call	☐ Pain limits washing / dressing: urgent review	☐ Acute severe illness: emergency help
Heart symptoms	☐ None	☐ New palpitations: contact team promptly	☐ Palpitations with dizziness: urgent assessment	☐ Chest pain or fainting: emergency assessment	☐ Severe chest pain, collapse or breathing difficulty: 112/999
Fever / chills	☐ No fever or chills	☐ New chills / feeling unwell: call	☐ Fever: use the written urgent plan now	☐ Fever with marked weakness: urgent assessment	☐ Collapse / confusion / breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.