SACTCHECK · TREATMENT PASSPORT

# Weekly cetuximab + head and neck radiotherapy

## A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Cetuximab starts one week before radiotherapy: a 400 mg/m² first infusion, then 250 mg/m² every week during radiation until radiotherapy ends. The number of weekly doses follows the actual radiotherapy plan.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

**Insert numbers here**

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

**Insert numbers here**

Ask for medical oncology

EVENINGS / OVERNIGHT

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00207/head_neck/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending

# Weekly cetuximab + head and neck radiotherapy

Cycle number: _____ Start date: _____ Next review: _____

Record actual treatment received, dose changes, tests and contact plans with your team. A stamp records a completed treatment; it does not authorise the next dose.

**Cetuximab before radiotherapy** · copy this sheet for another cycle of the same phase.

### Cetuximab · EGFR-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Eye symptoms** | ☐ None | ☐ New irritation: report | ☐ Pain, redness or light sensitivity: call promptly | ☐ Blurred vision / severe eye pain: urgent review | ☐ Sudden major vision loss: emergency assessment |
| **Breathing** | ☐ Usual breathing | ☐ New cough / change: call | ☐ Worsening breathlessness: urgent advice | ☐ Breathless at rest: urgent assessment | ☐ Severe breathing difficulty: 112/999 |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Muscle cramps** | ☐ None | ☐ Mild cramps: record | ☐ Repeated cramps: contact team | ☐ Severe pain / weakness or dark urine: urgent review | ☐ Sudden major weakness / collapse: 112/999 |
| **Infusion symptoms** | ☐ None | ☐ New flushing, chills or itching: tell the nurse immediately | ☐ Symptoms during infusion: tell the nurse immediately | ☐ Wheeze, swelling or faintness: immediate urgent help | ☐ Severe breathing difficulty or collapse: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Weekly cetuximab + head and neck radiotherapy

Cycle number: _____ Start date: _____ Next review: _____

Record actual treatment received, dose changes, tests and contact plans with your team. A stamp records a completed treatment; it does not authorise the next dose.

**Weekly cetuximab with radiotherapy** · copy this sheet for another cycle of the same phase.

## Cetuximab · EGFR-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Eye symptoms** | ☐ None | ☐ New irritation: report | ☐ Pain, redness or light sensitivity: call promptly | ☐ Blurred vision / severe eye pain: urgent review | ☐ Sudden major vision loss: emergency assessment |
| **Breathing** | ☐ Usual breathing | ☐ New cough / change: call | ☐ Worsening breathlessness: urgent advice | ☐ Breathless at rest: urgent assessment | ☐ Severe breathing difficulty: 112/999 |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Muscle cramps** | ☐ None | ☐ Mild cramps: record | ☐ Repeated cramps: contact team | ☐ Severe pain / weakness or dark urine: urgent review | ☐ Sudden major weakness / collapse: 112/999 |
| **Infusion symptoms** | ☐ None | ☐ New flushing, chills or itching: tell the nurse immediately | ☐ Symptoms during infusion: tell the nurse immediately | ☐ Wheeze, swelling or faintness: immediate urgent help | ☐ Severe breathing difficulty or collapse: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

## Radiotherapy · Local treatment

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Mouth soreness / ulcers** | ☐ None | ☐ Mild soreness; eating normally | ☐ Painful; soft food needed: call | ☐ Cannot eat / drink enough: urgent review | ☐ Major breathing difficulty: 112/999 |
| **Taste / appetite** | ☐ Usual eating | ☐ Taste change: record | ☐ Eating less / weight falling: contact team | ☐ Unable to eat or drink enough: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Weight** | ☐ Usual weight | ☐ Small change: record | ☐ Ongoing loss / swelling: contact team | ☐ Rapid change with breathlessness: urgent review | ☐ Severe breathing difficulty / collapse: 112/999 |
| **Skin** | ☐ No change | ☐ New small change: report | ☐ Persistent rash / itch: contact team | ☐ Painful, spreading or blistering rash: urgent review | ☐ Severe skin reaction with acute illness: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.