SACTCHECK · TREATMENT PASSPORT

Paclitaxel + cetuximab every 4 weeks

A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Paclitaxel is given on Days 1, 8 and 15, with cetuximab on Days 1 and 15 of each 28-day cycle. There is no scheduled treatment dose on Day 22. Both medicines continue while appropriate, until progression or unacceptable side effects; the protocol does not set a fixed total number of cycles.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00696/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending

Paclitaxel + cetuximab every 4 weeks

Cycle number: _____ Start date: _____ Next review: _____

Record the date, medicines actually received, delays or omissions, test results and concerns with your team. Use a separate record for any changed treatment phase.

Combination treatment · copy this sheet for another cycle of the same phase.

Cetuximab · EGFR-targeted antibody

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Skin / nail changes	☐ None	☐ Mild change: report	☐ Pain or daily tasks affected: call	☐ Spreading infection or blistering: urgent review	☐ Severe illness / collapse: emergency help
Eye symptoms	☐ None	☐ New irritation: report	☐ Pain, redness or light sensitivity: call promptly	☐ Blurred vision / severe eye pain: urgent review	☐ Sudden major vision loss: emergency assessment
Breathing	☐ Usual breathing	☐ New cough / change: call	☐ Worsening breathlessness: urgent advice	☐ Breathless at rest: urgent assessment	☐ Severe breathing difficulty: 112/999
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Muscle cramps	☐ None	☐ Mild cramps: record	☐ Repeated cramps: contact team	☐ Severe pain / weakness or dark urine: urgent review	☐ Sudden major weakness / collapse: 112/999
Infusion symptoms	☐ None	☐ New flushing, chills or itching: tell the nurse immediately	☐ Symptoms during infusion: tell the nurse immediately	☐ Wheeze, swelling or faintness: immediate urgent help	☐ Severe breathing difficulty or collapse: emergency help
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Paclitaxel · Taxane chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help
Fever / chills	☐ No fever or chills	☐ New chills / feeling unwell: call	☐ Fever: use the written urgent plan now	☐ Fever with marked weakness: urgent assessment	☐ Collapse / confusion / breathing difficulty: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999
Muscle / joint pain	☐ None	☐ Mild aches; normal activity	☐ Pain limits tasks / sleep: call	☐ Pain limits self-care: urgent review	☐ Sudden severe deterioration: emergency help
Skin / hair	☐ No change	☐ Dry skin / hair loss: record	☐ Persistent rash / skin change: contact team	☐ Painful, spreading or blistering rash: urgent review	☐ Severe skin reaction with acute illness: emergency help
Infusion symptoms	☐ None	☐ New flushing, chills or itching: tell the nurse immediately	☐ Symptoms during infusion: tell the nurse immediately	☐ Wheeze, swelling or faintness: immediate urgent help	☐ Severe breathing difficulty or collapse: emergency help
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Paclitaxel + cetuximab every 4 weeks

Cycle number: _____ Start date: _____ Next review: _____

Record the date, medicines actually received, delays or omissions, test results and concerns with your team. Use a separate record for any changed treatment phase.

Day 8 within the same 28-day cycle · copy this sheet for another cycle of the same phase.

Paclitaxel · Taxane chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help
Fever / chills	☐ No fever or chills	☐ New chills / feeling unwell: call	☐ Fever: use the written urgent plan now	☐ Fever with marked weakness: urgent assessment	☐ Collapse / confusion / breathing difficulty: 112/999
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999
Muscle / joint pain	☐ None	☐ Mild aches; normal activity	☐ Pain limits tasks / sleep: call	☐ Pain limits self-care: urgent review	☐ Sudden severe deterioration: emergency help
Skin / hair	☐ No change	☐ Dry skin / hair loss: record	☐ Persistent rash / skin change: contact team	☐ Painful, spreading or blistering rash: urgent review	☐ Severe skin reaction with acute illness: emergency help
Infusion symptoms	☐ None	☐ New flushing, chills or itching: tell the nurse immediately	☐ Symptoms during infusion: tell the nurse immediately	☐ Wheeze, swelling or faintness: immediate urgent help	☐ Severe breathing difficulty or collapse: emergency help
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.