SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Sunitinib 37.5 mg · pancreatic NET

Progressive unresectable or metastatic well-differentiated pancreatic neuroendocrine tumour under NCCP 00327a. · NCCP 00327 · version 5 · 28-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
Sunitinib is a daily targeted capsule used to help control a pancreatic NET that cannot be removed or has spread and is progressing. The team reviews control, side effects and your priorities.

Schedule
Take the prescribed sunitinib dose once daily continuously, at the same time, consistently with or without food. The 28-day cycle is a review period, with no planned capsule-free week.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Sunitinib
TARGETED THERAPY CAPSULE

- ☐ High blood pressure and heart dysfunction
- ☐ Painful hands and feet / skin changes
- ☐ Diarrhoea, mouth soreness, fatigue and poor appetite
- ☐ Low blood counts, infection and bleeding
- ☐ Thyroid dysfunction and blood-glucose changes
- ☐ Kidney injury / protein in urine and liver abnormalities
- ☐ Rare severe skin, neurological or vascular effects; impaired wound healing

Discuss the pancreatic NET indication, benefit and trial limitations, daily dosing, blood-pressure and heart risks, thyroid/glucose effects, hand-foot symptoms, bleeding, wound healing, interactions and the testing plan.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00327 v5.

SACTCheck v0.84.37
Developed by Dr Paul O'Brien