Sunitinib 37.5 mg · pancreatic NET

Daily targeted capsules to help control progressing pancreatic NET.

Progressive unresectable or metastatic well-differentiated pancreatic neuroendocrine tumour under NCCP 00327a. · NCCP 00327 v5 · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Progressive unresectable or metastatic well-differentiated pancreatic neuroendocrine tumour under NCCP 00327a.

My overall treatment

Once daily continuously; 28-day reviews, with no planned rest week.

Continuous course
Continue while benefit and tolerability support it; no fixed final cycle.

A typical 28-day cycle

Day 1–7	Day 8–28
Start with a clear prescription	Keep taking daily unless told to hold

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Bloods / BP / thyroid** Required by regimen	Counts, kidneys/liver, thyroid and BP; urine and cardiac assessment as indicated.
During	**Cycle bloods / BP** Required by regimen	Counts, kidneys/liver before each cycle; BP at every attendance.
Every 12 weeks	**Thyroid** Required by regimen	Additional urine, glucose or heart checks as indicated.
Review / after	**Response and recovery** Individual plan	Scans, NET markers and follow-up on your team’s plan.

Between treatments

- Report painful hands/feet, diarrhoea, bleeding, swelling or low-glucose symptoms early.
- Chest pain, severe breathlessness or severe headache with vision change needs emergency help.

Swallow whole, consistently with or without food. Missed dose: no extra capsule; take the usual prescribed dose next day.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital’s temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Head / thyroid / energy

Sunitinib

Thyroid changes can contribute to tiredness. Severe headache, seizure, confusion or visual change needs emergency assessment.

2 Mouth / jaw

Sunitinib

Report mouth soreness, eating problems or jaw pain. Discuss dental work with your team before treatment or procedures.

3 Heart / blood pressure

Sunitinib

Blood pressure may rise without symptoms. Keep checks; new swelling or breathlessness needs advice, and chest pain needs emergency care.

4 Blood / bleeding

Sunitinib

Low counts can increase infection or bleeding risk. Report fever, sudden illness, unusual bruising or bleeding; heavy bleeding is an emergency.

5 Liver

Sunitinib

Liver tests may change before symptoms appear. Attend blood tests and report yellow skin or eyes, dark urine or severe abdominal pain.

6 Bowel / fluids

Sunitinib

Diarrhoea can cause dehydration. Follow the written plan and call for persistent symptoms, dizziness or inability to drink.

7 Hands / feet / skin


Sunitinib

Report painful redness, swelling or blisters early. Severe widespread blistering or peeling requires urgent help.

8 Kidneys / glucose

Sunitinib

Urine protein and kidney or glucose changes may need tests. Report reduced urine, swelling, shakiness, sweating or confusion.



Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call urgently for severe diarrhoea or dehydration, bleeding, new swelling, much less urine, severe skin blistering or symptoms of low blood sugar. Chest pain, severe breathlessness, collapse, seizure or severe headache with vision changes needs emergency assessment.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Progressive unresectable or metastatic well-differentiated pancreatic neuroendocrine tumour under NCCP 00327a. · NCCP 00327 v5 · Document revised 5 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Sunitinib 37.5 mg · pancreatic NET

Once daily continuously; 28-day reviews, with no planned rest week.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling review record; no fixed final cycle.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.