NCCP 00726 · VERSION 3A

Pertuzumab + trastuzumab maintenance

Pertuzumab and trastuzumab are given by infusion every 21 days after the chemotherapy phase of the metastatic treatment plan. Duration depends on benefit, tolerance and the agreed plan; this is not a fixed one-year adjuvant course.

Draft — independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Hair / scalp

Whole treatment plan

Tell the team about changes affecting wellbeing or daily life.

2 Mouth

Treatment symptoms

Painful ulcers or difficulty eating and drinking need advice.

3 Breathing


HER2 treatment

New cough or breathlessness needs prompt assessment.

4 Heart / circulation

HER2 treatment

Fainting, chest pain or severe breathing difficulty: urgent help. Keep scheduled cardiac checks.



5 Bowel

Treatment medicines

Report diarrhoea early, especially with poor fluid intake, pain or fever.

6 Nerves / joints

Previous treatment

Report lasting numbness, painful joints or difficulty with daily activities.

7 Blood / infection

Whole treatment plan

Fever, shivers or suddenly unwell: contact oncology immediately.

8 Skin / liver symptoms

Treatment medicines

Severe rash, blistering or yellow skin/eyes needs prompt advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

NCCP 00726 · PRACTICAL SUPPORT

Your treatment, at home.

Important medicine-specific effects

Pertuzumab

- Diarrhoea and dehydration
- Infusion / allergic reaction
- Heart pumping-function reduction
- Rash and fatigue

Trastuzumab

- Heart pumping-function reduction / heart failure
- Infusion or allergic reaction
- Diarrhoea and fatigue
- Rare serious lung effects

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Infusion planning

Confirm the last treatment date with the team; long gaps can change the loading-dose plan.

Heart checks

Echoes continue on maintenance. Report new breathlessness, ankle swelling or rapid weight gain.

Bowel symptoms

Report diarrhoea early and follow the agreed hydration/support plan.

Response and daily life

Keep response-review appointments. Ask for support with fatigue, activity, sexual wellbeing and practical concerns.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

CUH · MON–FRI 8AM–5PM

087 181 9645

Acute Oncology Service

CUH · WEEKENDS / BANK HOLIDAYS 9AM–5PM

021 4922000

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00726/

Sources and review status are available online. Follow your own prescription and hospital instructions.