SACTCHECK TREATMENT ASSESSMENT

Pembrolizumab + Carboplatin/Paclitaxel -> AC

NCCP 00857 | 3
SACTCheck v0.83.7

TUMOUR GROUP	INDICATION	COURSE
Breast	Pembrolizumab in combination with chemotherapy as neoadjuvant treatment, then continued as monotherapy as adjuvant	Neoadjuvant

TREATMENT CONTEXT	ASSESSMENT ID	ASSESSMENT TIME
Not entered	JSON-DEMO-001	17/10/2026, 03:27

ENCODED CRITERIA RESULT

1 encoded criterion requires clinical review

Review the highlighted protocol criteria and the current NCCP source before the final treatment decision.

ENTERED VALUES AND ENCODED CRITERIA

Adjuvant pembrolizumab

DOMAIN	ENTERED VALUE	ENCODED CRITERION	RESULT
Pembrolizumab pneumonitis grade (source CTCAE)	Grade 3	>= 3	**Criterion not met - urgent review**

COMPONENT-SPECIFIC PROTOCOL ACTIONS

Protocol-level guidance; verify prescribed dose/current level

Pembrolizumab: Discontinue - Not to be administered
Permanently stop for grade 3-4 or recurrent grade 2 pneumonitis. page 8 | Table 4

UNASSESSED DOMAINS

ECOG performance status, HER2 status, Hormone-receptor status, Known hypersensitivity to a regimen medicine, Pregnant, Breastfeeding, History of ILD, Progression during platinum chemotherapy + 30 more

Blank domains were not assumed normal.

CLINICIAN DECISION

Not documented
Not documented

CLINICAL DECISION SUPPORT - NOT TREATMENT CLEARANCE

SACTCheck applies encoded NCCP protocol criteria to clinician-entered information. This output does not constitute treatment clearance and does not replace complete clinical assessment, the current official NCCP protocol, prescribing information, local policy or professional judgement. The responsible oncology clinician retains responsibility for the final treatment decision.