NCCP 00259 · VERSION 6

Vinorelbine capsules · advanced breast cancer

ONCE WEEKLY. First 3 administrations 60 mg/m², total maximum 120 mg. From administration 4,80 mg/m² (maximum 160 mg) only after the team reviews nadirs and toxicity.60 or 50 mg/m²/lower may be needed. Continue weekly while helpful and tolerated; no fixed total number of weeks.

Make treatment fit into daily life

Take with some food; swallow capsules whole with water. Do not chew, suck or dissolve. The exact total dose may require more than one capsule; pharmacy provides the strengths and number. Verify the weekly date, dose and capsule count each time. Do not take another dose if vomiting occurs within a few hours. Contact the team and record the dose taken. Do not swallow damaged capsules. Return them to pharmacy; immediately wash any contact with contents using water or saline. Missed/late dose: contact the team for the next date. Do not double or move the weekly schedule yourself. This is pragmatic medication-safety advice, not an extra NCCP catch-up schedule.

Visits and tests

Home dosing still needs weekly blood tests and explicit team clearance. Arrange a reliable weekly routine and a way to contact the team before taking a dose if unwell.

Treatment preparation

Record actual dose, capsule strengths/count, date taken, vomiting and side effects in the passport. An unfilled date is not a request for patients to calculate treatment dates.

At home

Ensure access to a refrigerator if required by the dispensed pack; discuss travel/transport with pharmacy.

Contacts / follow-up

Current Vinorelbine Accord SmPC: refrigerator 2–8°C. Confirm the dispensed brand/pack instructions; keep in original packaging and out of children’s reach. Return unused or damaged capsules to pharmacy; follow cytotoxic handling advice.

Who arranges each check?

Agree the due window, booking team, result reviewer and whom to contact if an appointment or result is missing. This reference page does not book or record a test.

Sources and review

The NCCP regimen supplies the treatment rules. Separate guidelines supply the longer-term monitoring recommendations, with their qualifications retained.

- NCCP 00259 v6 · official regimen
- Current Irish product information · safety supplement
- NCCP antiemetic guidance v6
- NCCP NCIS antiemetic medicines v8
- Freyer G et al. Journal of Clinical Oncology. 2003;21:35–40. DOI 10.1200/JCO.2003.09.057

Source reconciled; independent clinical, pharmacy and patient/PPI review pending