Vinorelbine IV · advanced breast cancer

30 mg/m² onDays 1 and 8 every 21 days. Consultant may consider 25 mg/m² for poor marrow reserve; the dose is individualised after toxicity. Repeated 21-day cycles until progression/unacceptable toxicity; team review determines continuation.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Blood counts and infection

Vinorelbine

White blood cells can fall, raising the risk of infection. Anaemia can cause tiredness or breathlessness; low platelets can cause bruising or bleeding. Blood tests guide delays and dose changes. Fever, shaking chills or feeling suddenly unwell need the oncology emergency contact

3 Tingling, numbness or weakness


Vinorelbine

Nerve effects may change sensation, grip, walking or bowel movement. Report changes early because the next dose may need to wait or be reduced.

5 Uncommon serious symptoms

Vinorelbine

New chest pain, sudden or worsening breathlessness, or severe allergic symptoms need urgent medical help. Previous heart disease and interacting medicines need review before treatment.



2 Constipation, nausea and mouth problems

Vinorelbine

The bowel can slow down and nausea, vomiting, diarrhoea or sore mouth can occur. Use the prescribed bowel and anti-sickness plan. Severe abdominal pain, swelling, repeated vomiting or inability to pass stool/gas needs urgent assessment for ileus: do not simply take extra

4 Fatigue, hair and liver tests

Vinorelbine

Tiredness, hair thinning/loss and liver-test changes can occur. Pace activities and tell the team if eating, drinking or daily tasks become difficult.

6 Infusion site

Vinorelbine

Tell staff immediately about burning, pain, swelling or leakage.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects are shown. Symptoms can occur on any day; this is not a complete list.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Vinorelbine

- Low blood counts, infection and bleeding
- Nausea, vomiting and fatigue
- Constipation or bowel obstruction
- Nerve effects and uncommon lung/ heart toxicity
- Infusion-site injury

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Sickness plan

No routine preventive anti-sickness medicine is usually required for this IV regimen. Ask for a rescue plan and report uncontrolled symptoms.

Bowel care

Agree a constipation plan, particularly if using opioids or ondansetron. Severe abdominal pain, swelling, repeated vomiting or inability to pass stool/gas needs urgent assessment; do not simply add laxatives.

Medicines and vaccines

Check every new prescription, supplement and vaccine with the team; avoid grapefruit. Live vaccines and interacting medicines need review.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00232/

Sources and review status are available online. Follow your own prescription and hospital instructions.