Capivasertib + fulvestrant

Capivasertib is taken twice daily for four days, then stopped for three days, each week. Fulvestrant injections are given on Days 1 and 15 of the first 28-day cycle, then Day 1 of later cycles. Treatment continues while helpful and tolerated.

Targeted tablets + endocrine injections · selected medicine-specific effects. Full details and sources are in the complete reference.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Blood sugar

Capivasertib

New thirst, passing more urine or marked tiredness: contact the team promptly.

2 Bowel / hydration

Capivasertib

Start the agreed diarrhoea plan; seek help if severe, persistent or you cannot drink.

3 Skin / mouth

Capivasertib

Report new rash or mouth soreness. Blistering or peeling needs urgent help.

4 Injection sites

Fulvestrant

Report bruising, unusual bleeding, radiating leg pain, numbness or weakness.

5 Hot flushes / joints


Fulvestrant

Support is available for flushes and joint or muscle symptoms.

6 Jaw / teeth

Capivasertib with bone medicines

Report jaw pain, loose teeth or a wound that does not heal, particularly with bone-modifying treatment.



Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Seek urgent assessment for unusual thirst or urination, severe diarrhoea, inability to drink, or a new rash. Vomiting with abdominal pain, breathing difficulty, fruity breath, confusion or marked sleepiness can signal ketoacidosis: seek emergency help without waiting for a high glucose reading. Blistering or peeling skin also needs urgent help. Severe chest pain, major breathing difficulty, collapse or stroke symptoms: 112 / 999.

YOUR PRACTICAL TREATMENT PLAN

Know the next step.

Taking treatment

Capivasertib is taken twice daily for four days, then stopped for three days, each week. Fulvestrant injections are given on Days 1 and 15 of the first 28-day cycle, then Day 1 of later cycles. Treatment continues while helpful and tolerated.

The usual starting capivasertib dose is 400 mg twice daily, about 12 hours apart, on the prescribed dosing days. Swallow tablets whole, with or without food; avoid grapefruit. Your prescription may use a lower dose. Each usual 500 mg fulvestrant dose is two slow injections into muscle, one in each buttock; no IV port or pump is required.

Checks and follow-up

Check blood count, kidney and liver function before treatment and each cycle. Check fasting glucose and HbA1c before starting. Fasting glucose is checked at weeks 1, 2, 4, 6 and 8, then monthly; ideally before the tablet dose on day 3 or 4 of a dosing week. HbA1c is checked every three months, with an extra week-4 check for diabetes, prediabetes or high baseline glucose. Diabetes, new high glucose or metformin initiation needs extra checks under a written plan.

The whole-treatment map shows the checks for this exact treatment setting.

Support between reviews

Glucose plan

Agree when to check glucose and when to call. Use your written diabetes-medicine plan on tablet-taking and tablet-free days; do not change existing diabetes medicines yourself.

Diarrhoea and fluids

Have the prescribed diarrhoea medicine available. Follow the agreed early-treatment instructions and contact the team for persistent symptoms, dizziness or reduced urine.

Skin and injections

Report rash early and use skin treatments as advised. Tell the injector about anticoagulants, bruising and any previous leg nerve symptoms.

Oncology daytime: Insert numbers here. **Out of hours:** Insert numbers here. Follow your own hospital’s urgent-care plan. Emergency: 112 / 999.

Your complete treatment reference

https://sactcheck.com/patient/00947/

Public information only. No personal information is encoded.