NCCP 00947 · VERSION 1

Capivasertib + fulvestrant

Capivasertib is taken twice daily for four days, then stopped for three days, each week. Fulvestrant injections are given on Days 1 and 15 of the first 28-day cycle, then Day 1 of later cycles. Treatment continues while helpful and tolerated.

Your treatment and review passport

Bring this to reviews. Record symptoms, treatment changes and questions when helpful. You do not need to complete a daily diary or calculate a toxicity grade.

The four sheets are reusable review periods, not a prescribed number of cycles. Follow the treatment and rest days agreed with your team; do not calculate a restart from this sheet.

These patient-language prompts are not a validated questionnaire or a formal CTCAE assessment. Urgent symptoms override the form.

Emergency: 112 / 999. Severe chest pain, major breathing difficulty, collapse or stroke symptoms need emergency help.

Contact your treating team promptly: Seek urgent assessment for unusual thirst or urination, severe diarrhoea, inability to drink, or a new rash. Vomiting with abdominal pain, breathing difficulty, fruity breath, confusion or marked sleepiness can signal ketoacidosis: seek emergency help without waiting for a high glucose reading. Blistering or peeling skin also needs urgent help.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your agreed out-of-hours pathway

Other hospitals: use the contacts your own team provides.

Your complete treatment reference

https://sactcheck.com/patient/00947/

Public information only. No personal information is encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

Capivasertib + fulvestrant

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Thirst / increased urination	☐	☐	☐	
Diarrhoea / dehydration	☐	☐	☐	
Rash / itching / mouth soreness	☐	☐	☐	
Nausea / appetite / fatigue	☐	☐	☐	
Injection pain / bleeding / nerve symptoms	☐	☐	☐	
Hot flushes / joint discomfort	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Seek urgent assessment for unusual thirst or urination, severe diarrhoea, inability to drink, or a new rash. Vomiting with abdominal pain, breathing difficulty, fruity breath, confusion or marked sleepiness can signal ketoacidosis: seek emergency help without waiting for a high glucose reading. Blistering or peeling skin also needs urgent help. Emergency symptoms: 112 / 999.

Capivasertib + fulvestrant

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Thirst / increased urination	☐	☐	☐	
Diarrhoea / dehydration	☐	☐	☐	
Rash / itching / mouth soreness	☐	☐	☐	
Nausea / appetite / fatigue	☐	☐	☐	
Injection pain / bleeding / nerve symptoms	☐	☐	☐	
Hot flushes / joint discomfort	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Seek urgent assessment for unusual thirst or urination, severe diarrhoea, inability to drink, or a new rash. Vomiting with abdominal pain, breathing difficulty, fruity breath, confusion or marked sleepiness can signal ketoacidosis: seek emergency help without waiting for a high glucose reading. Blistering or peeling skin also needs urgent help. Emergency symptoms: 112 / 999.

Capivasertib + fulvestrant

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Thirst / increased urination	☐	☐	☐	
Diarrhoea / dehydration	☐	☐	☐	
Rash / itching / mouth soreness	☐	☐	☐	
Nausea / appetite / fatigue	☐	☐	☐	
Injection pain / bleeding / nerve symptoms	☐	☐	☐	
Hot flushes / joint discomfort	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Seek urgent assessment for unusual thirst or urination, severe diarrhoea, inability to drink, or a new rash. Vomiting with abdominal pain, breathing difficulty, fruity breath, confusion or marked sleepiness can signal ketoacidosis: seek emergency help without waiting for a high glucose reading. Blistering or peeling skin also needs urgent help. Emergency symptoms: 112 / 999.

Capivasertib + fulvestrant

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Thirst / increased urination	☐	☐	☐	
Diarrhoea / dehydration	☐	☐	☐	
Rash / itching / mouth soreness	☐	☐	☐	
Nausea / appetite / fatigue	☐	☐	☐	
Injection pain / bleeding / nerve symptoms	☐	☐	☐	
Hot flushes / joint discomfort	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Seek urgent assessment for unusual thirst or urination, severe diarrhoea, inability to drink, or a new rash. Vomiting with abdominal pain, breathing difficulty, fruity breath, confusion or marked sleepiness can signal ketoacidosis: seek emergency help without waiting for a high glucose reading. Blistering or peeling skin also needs urgent help. Emergency symptoms: 112 / 999.