Docetaxel alone for previously treated non-small-cell lung cancer

To help control previously treated non-small-cell lung cancer and relieve symptoms.

Lung · previously treated advanced non-small-cell lung cancer (NSCLC) · NCCP 00203 v7 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Docetaxel by drip on Day 1, repeated every 21 days, for up to six cycles. Treatment may stop earlier if it is not helping or side effects are unacceptable. Your team confirms the prescribed course.

21 day view

Treatment timing in this window

Chemotherapy · Day 1 — Day 1

Before starting	Treatment every 21 days	After treatment
Confirm your blood tests, current medicines and steroid plan.	Day 1 drip, then time at home. Up to six cycles, with a review before each.	Review benefit, recovery and the next care plan with your team.

A typical 21-day cycle

Day 1	Day 2	Day 3–20	Day 21
Treatment day	Day after treatment	Between visits	Prepare for the next cycle

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before starting	**Blood count, kidneys and liver** Required by regimen	Before the first dose
During treatment	**Blood count, kidneys and liver** Required by regimen	Before every 21-day cycle
During / after	**Tumour markers, scans or extra tests** If clinically indicated	Scans: team-agreed timing. Markers: only if relevant; no routine schedule.
After treatment	**Recovery and follow-up** Team-agreed plan	Team agrees timing; follow ongoing symptoms or abnormal results

Between treatments

- Day 2: finish prescribed steroid. Between visits, watch for infection, bowel changes, numbness or swelling; timing varies.
- Before another cycle: blood tests and symptom review. Delays or early stopping may be needed.

Take the prescribed steroid from the day before each infusion. Tell the team if any doses were missed. Bring your medicine list and report symptoms since the last visit. Ask when your next blood tests and cancer review will be.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hands / feet

Docetaxel

Report numbness, weakness or difficulty walking or using your hands.

3 Hair / scalp

Docetaxel

Hair loss can occur; ask about support before treatment.

5 Mouth


Docetaxel

Report painful ulcers, difficulty drinking or poor intake.

7 Skin / swelling

Docetaxel

Report swelling; blistering or peeling skin needs urgent advice.



2 Blood / infection

Docetaxel

Fever, shivers or sudden illness need immediate contact.

4 Breathing

Docetaxel

New or worsening cough or breathlessness needs urgent assessment.

6 Tummy / bowel

Docetaxel

Severe tummy pain or rapidly worsening diarrhoea needs urgent assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call the oncology contact number immediately for fever, shivers or sudden illness, severe or new breathing difficulty, chest pain, fainting, severe abdominal pain or persistent diarrhoea. Tell staff immediately about infusion symptoms. Report new numbness, weakness, painful skin changes, swelling or visual changes promptly; do not wait for the next cycle. Severe tummy pain, rapidly worsening diarrhoea, blistering or peeling skin, or new breathing symptoms also need urgent assessment.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Lung · previously treated advanced NSCLC · NCCP 00203 v7 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Docetaxel

Day 1 every 21 days; up to 6 cycles

Phase: _____ Planned cycles: 6

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Team stamps after each completed cycle.

C1	C2	C3
Date / initials	Date / initials	Date / initials
C4	C5	C6
Date / initials	Date / initials	Date / initials

My treatment information

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.