Carboplatin · 21-day breast schedule

Carboplatin is infused on Day 1 of each 21-day cycle. The team selects the dose using the prescribed AUC and kidney function. Treatment continues while helpful and tolerated; there is no fixed breast course length.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Blood / infection

Carboplatin

Fever, unusual bruising or bleeding needs urgent advice.

2 Kidneys / urine

Carboplatin

Keep kidney tests and report reduced urine or swelling.

3 Hearing

Carboplatin

Report ringing in the ears or hearing changes.

4 Hands / feet

Carboplatin

Report tingling, numbness or altered balance early.

5 Nausea / appetite


Carboplatin

Follow the prevention plan; call if vomiting prevents fluids.

6 Allergic reaction

Carboplatin

During an infusion, report rash, chest tightness or breathlessness immediately.



Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects are shown. Symptoms can occur on any day; this is not a complete list.

Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Carboplatin

- Low blood counts, infection and bleeding
- Nausea, vomiting and fatigue
- Kidney, hearing and nerve effects
- Infusion allergy, particularly with repeat platinum exposure
- Fertility effects and harm to an unborn baby

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Preventing sickness

Take the prescribed anti-sickness medicines before treatment and on the following days, even if you feel well. Follow your own prescription and contact the team if nausea or vomiting is not controlled.

Blood counts

Keep blood tests and use the emergency contact for fever, chills or bleeding. Preventive growth-factor injections depend on individual risk.

Kidneys and other medicines

Tell the team about every medicine or supplement, especially treatments that affect kidneys, hearing or bleeding. Follow individual drinking advice.

Steroid and rescue plan

The anti-sickness plan may affect sleep, glucose and bowel symptoms. Confirm the exact doses and last day of each medicine; contact the team if symptoms persist.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00261/

Sources and review status are available online. Follow your own prescription and hospital instructions.