Cisplatin + etoposide with radiotherapy

Chemotherapy and radiation work together to treat limited-stage small-cell lung cancer. Your team explains your treatment aim and choices.

Limited-stage small-cell lung cancer · combined chemotherapy and radiation · NCCP 00279 v8 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

For limited-stage lung cancer: intravenous (IV) cisplatin/etoposide every 21 days. Oral, split-dose, carboplatin or 28-day plans need personalised information.

My overall treatment

Reference IV plan: four 21-day cycles. Day 1: cisplatin + etoposide. Days 2–3: etoposide. Radiation follows its own appointments. Capsules, split cisplatin, carboplatin substitution or 28-day cycles require a different personalised prescription.



Chemotherapy

Three treatment days per reference cycle, then recovery. Your team confirms each cycle after safety checks.

Radiotherapy

Usually starts with cycle 1. Radiation oncology sets the dose and visits; chemotherapy delays need separate coordination.

After treatment

Your team reviews recovery, persistent symptoms and response. Scan or marker timing is individual.

A typical 21-day cycle

Day 1	Day 2–3	Day 4–7	Day 8–21
reference combination	Days 2–3: etoposide	Support and recovery	Between chemotherapy cycles

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Blood count, kidneys and liver** Required by regimen	Before treatment
Before	**Audiology / creatinine clearance** If clinically indicated	When clinically indicated
During	**Bloods, hearing and nerves** Required by regimen	Before every chemotherapy cycle
After / review	**Recovery and response** Team-agreed plan	Team-agreed timing; no universal interval

Between treatments

- Follow the prescribed anti-sickness course and fluid advice; call early if eating or drinking becomes difficult.
- Fever, new breathlessness, uncontrolled vomiting or inability to drink needs urgent advice, including after treatment ends.

Confirm your prescription, radiation appointments and blood tests. Bring your medicine list and daytime / overnight contact numbers.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hearing / nerves


Cisplatin

Report ringing, reduced hearing, numbness or weakness before the next dose.

2 Mouth / swallowing

Treatment and chest radiation

Soreness may affect food and fluids. Contact the team early; inability to drink is urgent.



3 Breathing

Chemoradiation

New or worsening breathlessness needs urgent assessment, including after treatment finishes.

4 Sickness / appetite

Chemotherapy

Follow the anti-sickness plan. Call if vomiting continues or fluids will not stay down.

5 Kidneys / fluid balance

Cisplatin

Blood tests and fluids protect kidneys. Report much less urine or poor intake.

6 Blood cells / IV site

Chemotherapy

Fever or sudden illness can be serious. Report bleeding or infusion pain promptly.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call oncology immediately for fever, shaking chills, sudden illness, new breathlessness, uncontrolled vomiting, inability to swallow fluids, much less urine or unusual bleeding. Severe breathing difficulty, chest pain, collapse or heavy bleeding needs emergency care. Tell staff immediately about infusion pain, swelling or allergy symptoms.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Limited-stage small-cell lung cancer · combined chemotherapy and radiation · NCCP 00279 v8 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Cisplatin + etoposide / RT

Day 1: both drugs. Days 2–3: etoposide. Usually every 21 days ×4; RT plan separate.

Phase: _____ Planned cycles: 4

Daytime (EXAMPLE) 087 12345678
Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Team stamps after each completed cycle.

C1 Date / initials

C2 Date / initials

C3 Date / initials

C4 Date / initials

My treatment information

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.