INFORMATION TO KEEP WITH YOU

# My complete treatment handbook

## Pertuzumab + trastuzumab · with selected chemotherapy

Understand your treatment. Know what to expect. Keep your plan to hand.

**Displayed calendar**
**21 days**
See the active treatment phase

**Treatment stages**
**2**
Explained in the treatment reference

**Regimen source**
**NCCP 00350**
Version 3

## Treatment stages

01 **Before surgery · selected chemotherapy + antibodies**
21-day cycle

02 **After surgery · agreed adjuvant plan**
21-day cycle

Your team confirms the active stage and dates. The full treatment guide explains each step.

## Inside your handbook


| | | PAGE |
|---|---|---|
| 01 | Understanding your treatment | 2 |
| 02 | Your schedule, tests and appointments | 5 |
| 03 | Symptoms and supportive care | 12 |
| 04 | Your record and contact card | 17 |
| 05 | Sources and further information | 20 |


**My name** _____

**Hospital / oncology team** _____

Use the urgent instructions and contact numbers in your treatment guide. Your team must replace example numbers before patient use. Your individual treatment plan takes precedence.

CHAPTER 01

# Understanding your treatment

IN THIS CHAPTER

Why this treatment is offered, what its medicines do, and how the agreed plan fits your care.

## When to get help

**Emergency: call 112 / 999**

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

**Contact your oncology service immediately**

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Report new breathlessness, chest symptoms, severe diarrhoea, dehydration, jaundice, unusual bleeding or sudden illness promptly. Do not wait for the next planned test.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

## What each medicine does

Treat HER2-positive early breast cancer as part of a plan intended to reduce the risk of it returning.

HER2-TARGETED ANTIBODY

### Pertuzumab

HER2-targeted antibody



**Blocks HER2 pairing with other receptors**

Pertuzumab binds a different area of HER2 from trastuzumab. It limits receptor pairing, so the two antibodies act together at different points on the same growth pathway.

HER2 pairing → Pertuzumab blocks pairing → Growth signals reduced

**What to watch for**

Keep heart-scan appointments. Report diarrhoea, new breathlessness, swelling or a treatment reaction.

HER2-TARGETED ANTIBODY

### Trastuzumab

HER2-targeted antibody



**Targets HER2 growth signals**

Trastuzumab binds to the HER2 protein on cells and helps reduce HER2-driven growth and recruit immune activity. HER2 testing identifies the cancers for which this approach is used.

HER2 on cell surface → Antibody binds → Growth signalling reduced

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

## Why this treatment?

Two HER2-targeted antibodies may be combined with selected chemotherapy for higher-risk HER2-positive early breast cancer. Cancer extent, lymph nodes, hormone receptors, heart function and the chosen chemotherapy influence the plan. This card explains the antibody component; the companion chemotherapy needs its own regimen guide and assessment.

**What could change the choice?**

Ask which cancer features, earlier treatments and health factors led to this recommendation, which alternatives were considered and when the decision will be reviewed.

## Your treatment and the next step

**Schedule**

intravenous (IV) pertuzumab and IV trastuzumab every 21 days. Before surgery: 3–6 cycles alongside the selected chemotherapy. After surgery: up to 18 cycles overall under the agreed adjuvant plan. If an anthracycline is used first, finish it before starting these antibodies with the taxane phase. Confirm the exact companion regimen and local access.

**After this phase**

The chosen chemotherapy, indication, response, tolerance and local access determine the next step. Do not treat this antibody framework as a complete chemotherapy protocol.

**Confirm the selected plan**

The source lists adjuvant pertuzumab reimbursement as not approved. Your team must confirm current access and the exact companion chemotherapy before this pathway is selected.

CHAPTER 02

# Your schedule, tests and appointments

IN THIS CHAPTER

Your treatment stages, checks and practical preparations, from before treatment through follow-up.

## The phases and how they connect

### Treatment stages

01 **Before surgery · selected chemotherapy + antibodies**
21-day cycle

02 **After surgery · agreed adjuvant plan**
21-day cycle

Your team confirms the active stage and dates. The full treatment guide explains each step.

This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

21-DAY CYCLE
### Before surgery · selected chemotherapy + antibodies

21-day cycle

Three to six antibody cycles with the selected chemotherapy. Anthracyclines, if used, finish before antibodies start with the taxane.

21-DAY CYCLE
### After surgery · agreed adjuvant plan

21-day cycle

Up to 18 cycles overall under the source; confirm individual plan and current access. Chemotherapy is separately specified.

Relative sequence, not equal lengths or booked dates. Your team confirms actual appointments and the active medicines.

BASELINE → TREATMENT → FOLLOW-UP

## Tests and milestones through treatment

Columns show sequence, not equal durations or appointments. A test is only complete when performed and reviewed. Echo timing and cancer-imaging timing are different. Symptoms can bring any check forward.

| Check | Select indication / chemotherapy | Anthracycline first, if used | Antibodies with taxane | Surgery / adjuvant continuation | Completion / follow-up |
|---|---|---|---|---|---|
| **Blood tests** | Baseline antibody and companion-regimen checks | Follow the chosen anthracycline source | Before each cycle plus companion-regimen checks | Follow the current medicines | Follow abnormal results / next plan |

| Check | Select indication / chemotherapy | Anthracycline first, if used | Antibodies with taxane | Surgery / adjuvant continuation | Completion / follow-up |
|---|---|---|---|---|---|
| **Echo / cardiac assessment** | Cardiac risk and baseline review | Reassess after anthracycline, before first HER2 dose | First echo 12 weeks after HER2 begins | Every 12 weeks during HER2; at completion | Follow-up for risk / chosen medicine |
| **Cancer imaging / markers** | Baseline disease assessment | Clinical response; imaging if concern | Response review as planned; not tied to echo | Reassess disease and treatment plan | Breast surveillance after early treatment |
| **Surgery / radiotherapy** | Agree indication and sequence | No automatic concurrent radiation | Coordinate with both teams | Local treatment as indicated | Review recovery and late effects |
| **Endocrine / bone health** | Check hormone receptors / fracture risk | Plan endocrine therapy if indicated | dual-energy X-ray absorptiometry (bone density) (DEXA) when relevant to bone-risk plan | Endocrine timing by the treating team | Individual DEXA / bone follow-up |

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00350 v3 · official regimen · ESMO 2024 early breast cancer guideline · treatment and follow-up · ESC 2022 cardio-oncology guideline · cardiac surveillance and risk · NICE NG101 · bone health and follow-up · ESMO metastatic breast cancer guideline · 2026

THE WHOLE TREATMENT JOURNEY

# Pertuzumab + trastuzumab · with selected chemotherapy: the whole pathway

## HER2-positive early breast cancer · neoadjuvant or adjuvant framework

**Read from left to right.** Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

**Treatment, tests and reviews across the course**

| Care / check | Select indication / chemotherapy | Anthracycline first, if used | Antibodies with taxane | Surgery / adjuvant continuation | Completion / follow-up |
|---|---|---|---|---|---|
| **Cancer medicines** NCCP | Confirm diagnosis and exact regimen | See the dose-day map: antibodies with selected chemotherapy | Start antibodies with the taxane | Confirm indication, companion and continuation | Review benefit / tolerance and next plan |
| **Surgery / radiotherapy** EARLY | Agree indication and sequence | No automatic concurrent radiation | Coordinate with both teams | Local treatment as indicated | Review recovery and late effects |
| **Blood tests** NCCP | Baseline antibody and companion-regimen checks | Follow the chosen anthracycline source | Before each cycle plus companion-regimen checks | Follow the current medicines | Follow abnormal results / next plan |
| **Echo / cardiac checks** NCCP · ESC | Cardiac risk and baseline review | Reassess after anthracycline, before first HER2 dose | First echo 12 weeks after HER2 begins | Every 12 weeks during HER2; at completion | Follow-up for risk / chosen medicine |
| **Scans / tumour markers** EARLY | Baseline disease assessment | Clinical response; imaging if concern | Response review as planned; not tied to echo | Reassess disease and treatment plan | Breast surveillance after early treatment |
| **Endocrine / DEXA** BONE | Check hormone receptors / fracture risk | Plan endocrine therapy if indicated | DEXA when relevant to bone-risk plan | Endocrine timing by the treating team | Individual DEXA / bone follow-up |

## What each check is for

1

### Blood tests

full blood count (FBC), kidney and liver profile at baseline and before every 21-day cycle. The team reviews the necessary results before a planned dose. The selected chemotherapy can require extra or more frequent checks.

Check safety before the next dose.

**Ask / act:** Agree the booking owner, result reviewer and action if overdue.

2

### Heart tests

Check heart pumping function before starting HER2 treatment. Repeat echo or multigated acquisition (heart pumping) scan (MUGA) every 12 weeks: around cycle 5, then cycle 9 if 21-day cycles stay on time. Earlier checks are needed for symptoms or abnormal findings. Count elapsed time from the first HER2 dose; changing phase or route does not restart the clock.

Monitor pumping function and relevant rhythm risk.

**Ask** / **act:** Agree the booking owner, result reviewer and action if overdue.

3



### Cancer imaging

Baseline breast and nodal imaging, with staging tests appropriate to the diagnosis. Review response during preoperative treatment and before surgery. After local treatment, arrange breast surveillance; routine computed tomography (CT), positron emission tomography (PET) or tumour-marker testing is not the usual screening plan for someone without symptoms.

Assess response or follow-up needs.

**Ask** / **act:** Agree the booking owner, result reviewer and action if overdue.

4

### Bone health

If the cancer is also hormone-receptor positive, endocrine treatment may form part of the wider plan. Bone-density testing is linked to aromatase inhibitors, ovarian suppression and fracture risk; it is not automatically required because a HER2 medicine is prescribed. Agree baseline DEXA where indicated and individual follow-up.

Assess fracture risk where relevant.

**Ask** / **act:** Agree the booking owner, result reviewer and action if overdue.

**At the next review, ask**

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

## Your practical treatment plan



STEP 1

### Before the first visit

Confirm the active regimen, blood results, cardiac checks, medicines review, route and consent discussion.



STEP 2

### Plan venous access

A cannula, peripherally inserted central catheter (PICC) or implanted port may be used. If a port is chosen, a trained team inserts it in a separate procedure with local anaesthetic and sometimes sedation; the team explains fasting, blood-thinner and wound instructions.

STEP 3

### Infusion day

Staff check the line, give prescribed premedicines and treatment, and observe for reactions. Tell them immediately about burning, pain, rash or breathlessness.

STEP 4

### Home and next review

This regimen does not routinely need a take-home infusion pump. Staff disconnect the infusion; a port is de-accessed when appropriate. Keep line-care and next-visit instructions.

#### Port insertion and removal

The line team explains placement, dressing care and when normal activities can resume. A port is accessed with a special needle during treatment. Removal of the implanted device is a separate planned procedure when no longer needed; do not remove or adjust it yourself.

#### PICC care and urgent line symptoms

A PICC usually needs regular dressing and flushing appointments even between treatments. Fever, redness, swelling, pain, leakage or damage requires prompt advice. Your hospital supplies the exact care instructions.

#### Visit time and transport

First visits can take longer because of loading doses, premedicines and observation. Ask the unit about expected time, transport and any escort needed after sedation or line insertion.

#### Phase-specific visit pattern

Infusion and observation times differ by drug and first versus later doses; the unit confirms the full visit duration, including blood review and preparation.

## When scans, heart checks and bloods happen

Agree who books each test, who reviews the result and what happens if a check is overdue.

### Blood tests

FBC, kidney and liver profile at baseline and before every 21-day cycle. The team reviews the necessary results before a planned dose. The selected chemotherapy can require extra or more frequent checks.

### Heart ultrasound and electrocardiogram (ECG)

Check heart pumping function before starting HER2 treatment. Repeat echo or MUGA every 12 weeks: around cycle 5, then cycle 9 if 21-day cycles stay on time. Earlier checks are needed for symptoms or abnormal findings. Count elapsed time from the first HER2 dose; changing phase or route does not restart the clock.

### Scans and tumour markers

Baseline breast and nodal imaging, with staging tests appropriate to the diagnosis. Review response during preoperative treatment and before surgery. After local treatment, arrange breast surveillance; routine CT, PET or tumour-marker testing is not the usual screening plan for someone without symptoms.

### Bone health and endocrine treatment

If the cancer is also hormone-receptor positive, endocrine treatment may form part of the wider plan. Bone-density testing is linked to aromatase inhibitors, ovarian suppression and fracture risk; it is not automatically required because a HER2 medicine is prescribed. Agree baseline DEXA where indicated and individual follow-up.

## How surgery and radiotherapy fit

The selected indication sets the order: neoadjuvant treatment → surgery → selected postoperative treatment; adjuvant treatment starts after surgery. Finish any anthracycline phase before starting these antibodies with the taxane. Radiation timing is agreed with the treating teams.

CHAPTER 03

# Symptoms and supportive care

IN THIS CHAPTER

Recognising changes, knowing when to call, and using your prescribed supportive-care plan.

## Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

| Treatment or check | Timing in this view |
|---|---|
| **Antibodies · Day 1 every 21 days** |  **Day 1** Selected chemotherapy has its own calendar |
| **Heart / breathing symptoms** |  **Days 1–21** Report at any time |
| **Diarrhoea** |  **Days 1–21** Report early |

**Planned treatment:** IV pertuzumab and IV trastuzumab every 21 days. Before surgery: 3–6 cycles alongside the selected chemotherapy. After surgery: up to 18 cycles overall under the agreed adjuvant plan. If an anthracycline is used first, finish it before starting these antibodies with the taxane phase. Confirm the exact companion regimen and local access.

**Next phase:** The chosen chemotherapy, indication, response, tolerance and local access determine the next step. Do not treat this antibody framework as a complete chemotherapy protocol.

**Heart checks:** Check heart pumping function before starting HER2 treatment. Repeat echo or MUGA every 12 weeks: around cycle 5, then cycle 9 if 21-day cycles stay on time. Earlier checks are needed for symptoms or abnormal findings. Count elapsed time from the first HER2 dose; changing phase or route does not restart the clock.

> **Why these bands stay broad:** Treatment markers show the planned schedule. Symptom bands are broad possibilities, not measured risk curves; serious symptoms can occur at any time. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00350 v3 · official regimen ↗

> **There are no guaranteed "safe days".** Serious symptoms need assessment on any day, even if the chart shows a usual recovery period.

### Between-treatment notes

### Day 1–21: The current treatment cycle

IV pertuzumab and IV trastuzumab every 21 days. Before surgery: 3–6 cycles alongside the selected chemotherapy. After surgery: up to 18 cycles overall under the agreed adjuvant plan. If an anthracycline is used first, finish it before starting these antibodies with the taxane phase. Confirm the exact companion regimen and local access. Your team confirms treatment dates and checks. Serious symptoms need advice on any day.

## Support around the medicines

### 1 Choose the companion chemotherapy

Confirm the exact chemotherapy regimen before issuing its anti-sickness or other supportive prescription. The minimal emetic risk of the two antibodies does not describe the whole combination.

### 2 Keep heart checks on the plan

An echo every 12 weeks is separate from blood tests or cancer imaging. Ask who books and reviews it.

### 3 Plan for symptoms

Keep your agreed contact numbers and instructions. New symptoms can need assessment before a routine appointment.

## Questions worth asking

**Who arranges my tests?**

Ask the team who books each test, who checks the result and who to contact if an appointment is overdue. This map does not book appointments.

**What if I become unwell between visits?**

Use the urgent contact plan for sudden illness, fever, chest symptoms, severe diarrhoea or poor fluid intake. Do not wait for a scheduled blood test or scan.

**Try explaining the plan back:** Which medicines are active now, which days do you receive or take them, and who is booking and reviewing your next tests?

### Pertuzumab

- Heart pumping-function reduction
- Diarrhoea and rash
- Fatigue

---

**Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

## Combination visit · select the chemotherapy

**Active medicines:** Pertuzumab + Trastuzumab + Separately selected chemotherapy

**Emetogenic risk:** Determined by the selected chemotherapy

Chemotherapy has not been specified here. A complete supportive prescription cannot be generated from this antibody framework alone.

### Additional phase-specific support

- Open the exact companion NCCP regimen before prescribing. Its antiemetics, premedication, granulocyte colony-stimulating factor (white-cell growth support) (G-CSF), blood thresholds and phase schedule apply.
- Antibody-specific premedication is not usually needed; consider paracetamol/antihistamine for the reaction history.
- If an anthracycline phase is selected, finish it before starting these antibodies with the taxane.
- Confirm indication and current access: NCCP00350b marks adjuvant pertuzumab reimbursement as not approved.
- Clinical template for individual review, not an issued prescription. Confirm allergy, pregnancy precautions, interactions and local policy.

**Before signing:** verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

---

## Antibodies only · if separately prescribed

**Active medicines:** Pertuzumab + Trastuzumab

**Emetogenic risk:** Minimal for antibodies only

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

### Additional phase-specific support

- Confirm that chemotherapy has actually ended or is separately held; do not apply this minimal-risk script to a combination visit.
- No routine cytotoxic antiemetic, taxane premedication or chemotherapy granulocyte colony-stimulating factor (white-cell growth support) (G-CSF) is carried forward.
- Continue cardiac checks every 12 weeks and at completion; a phase change does not reset the clock.
- Clinical template for individual review, not an issued prescription. Confirm allergy, pregnancy precautions, interactions and local policy.

**Before signing:** verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

# Pertuzumab + trastuzumab · with selected chemotherapy

intravenous (IV) pertuzumab and IV trastuzumab every 21 days. Before surgery: 3–6 cycles alongside the selected chemotherapy. After surgery: up to 18 cycles overall under the agreed adjuvant plan. If an anthracycline is used first, finish it before starting these antibodies with the taxane phase. Confirm the exact companion regimen and local access.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

## A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

### 1 Heart / breathing


**Pertuzumab + trastuzumab**

Keep echo appointments; report breathlessness, swelling or reduced exercise tolerance.

### 2 Diarrhoea

**Especially pertuzumab**

Report watery stools early, especially with dizziness or poor intake.




### 3 Infusion reaction

**Either antibody**

Tell staff immediately about rash, chest tightness or breathing difficulty.

### 4 Rash / skin

**HER2 antibodies**

Report persistent or severe skin changes.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

This map covers the two HER2 antibodies only. The selected chemotherapy has additional risks and requires its own patient guide.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

CHAPTER 04

# Your record and contact card

IN THIS CHAPTER

Space for your own cycle or review record, questions and notes, followed by your treatment card.

Record each cycle separately. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Use a new record sheet for each cycle or review. Copy or print another sheet when needed; the number of sheets does not prescribe the length of treatment.

Cycle number: _____ Start date: _____ Next review: _____

**Current regimen · reusable cycle record** · copy this sheet for another cycle of the same phase.

## Pertuzumab · HER2-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/ day or at night: call | □ 7+ extra stools/ day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

## Trastuzumab · HER2-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/ day or at night: call | □ 7+ extra stools/ day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Carry your treatment details

Print, cut out and fold. Follow the contact plan from your own hospital.

SACTCHECK · SACT TREATMENT

## Pertuzumab + trastuzumab · with selected chemotherapy

NCCP 00350 v3 · 21-day cycle

IV pertuzumab and IV trastuzumab every 21 days. Before surgery: 3–6 cycles alongside the selected chemotherapy. After surgery: up to 18 cycles overall under the agreed adjuvant plan. If an anthracycline is used first, finish it before starting these antibodies with the taxane phase. Confirm the exact companion regimen and local access.

| 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|

**Your contact plan**

Oncology team: Insert numbers here
Out of hours: Insert numbers here.

Evenings / overnight: instructed out-of-hours pathway or nearest ED.
**Emergency: 112 / 999.**

Other hospital / own contact: _____

QR: https://sactcheck.com/patient/00350/ · no patient information.

Draft for review · SACTCheck v0.84.16

CHAPTER 05

# Sources and further information

## IN THIS CHAPTER

The references behind this information and addresses you can use to find further help.

Use this information alongside the plan agreed with your treatment team. Your team confirms your medicines, doses and appointments.


1. **National Cancer Control Programme (NCCP) 00350 v3 · official regimen**
   https://healthservice.hse.ie/documents/6813/350_v3_Pertuzumab_Trastuz_Chemo.pdf
2. **ESMO 2024 early breast cancer guideline · treatment and follow-up**
   https://www.annalsofoncology.org/article/S0923-7534(23)05104-9/fulltext
3. **ESC 2022 cardio-oncology guideline · cardiac surveillance and risk**
   https://academic.oup.com/eurheartj/article/43/41/4229/6673995
4. **NICE NG101 · bone health and follow-up**
   https://www.nice.org.uk/guidance/ng101/chapter/Recommendations
5. **ESMO metastatic breast cancer guideline · 2026**
   https://www.annalsofoncology.org/article/S0923-7534(26)00902-6/fulltext
6. **EMA Herceptin product information**
   https://www.ema.europa.eu/en/documents/product-information/herceptin-epar-product-information_en.pdf
7. **eviQ · implanted port patient information**
   https://www.eviq.org.au/p/507
8. **eviQ · peripherally inserted central catheter (PICC) patient information**
   https://www.eviq.org.au/p/543
9. **HSE · chemotherapy side effects and urgent advice**
   https://www2.hse.ie/conditions/chemotherapy/side-effects/
10. **NOGG · section 7, aromatase inhibitors and fracture risk**
    https://www.nogg.org.uk/full-guideline/section-7-strategies-management-osteoporosis-and-fracture-risk
11. **NCCP NCIS antiemetics v8 (2025), Table 1**
    https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf
12. **ASCO · biomarkers in metastatic breast cancer**
    https://ascopubs.org/doi/10.1200/JCO.2015.61.1459
13. **EMA Perjeta product information**
    https://www.ema.europa.eu/en/documents/product-information/perjeta-epar-product-information_en.pdf