Weekly cisplatin + head and neck radiotherapy

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Cisplatin with radiotherapy

Active medicines: Cisplatin

Emetogenic risk: high

Medicine	Dose / route	When
High-emetic-risk anti-sickness plan	Clinician/pharmacy to complete	Before platinum and for delayed symptoms as prescribed Use the current local/NCCP policy; reconcile prevention and rescue medicines into one schedule.
Cisplatin hydration and electrolyte plan	Clinician/pharmacy to complete	Before and after cisplatin; home fluids individualised Check kidney function, fluid balance and relevant comorbidities; do not issue an automated fluid prescription.

Additional phase-specific support

- Ask for one schedule showing prevention medicines, rescue treatment and what to do if symptoms persist. Do not add, double or restart medicines from an old prescription.
- Cisplatin needs prescribed fluids before and after infusion. Follow the team's drinking advice and report much less urine, vomiting or inability to drink. Do not change fluid intake or diuretics without advice.
- A dietitian can tailor food texture, supplements and calorie or protein needs. Tell the team early if eating or drinking becomes difficult; do not wait for substantial weight loss. Fluid targets depend on kidney and heart health.
- Use the mouth-care plan provided by the team and tell them about ulcers, thrush, bleeding or pain. Inform your dentist about cancer treatment. Before any dental procedure, confirm timing with the oncology team.
- A speech and language therapist can assess swallowing and advise exercises or safer food and drink textures. Follow the individual assessment; a general guide cannot decide which textures are safe for you.
- Discuss fertility and pregnancy precautions before treatment and the medicine-specific duration of contraception afterwards. Ask about breastfeeding. Bring a full medicine, supplement and allergy list; check before adding over-the-counter or herbal products.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00385 v5 · pSee regimen treatment, tests and supportive-care sections · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-11-07 · SACTCheck v0.83.8 · Regimen-level drafts; prescriber and oncology-pharmacy verification required