Pembrolizumab 200 mg · colorectal

National Cancer Control Programme (NCCP) 00455 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Appointments

Pembrolizumab 200 mg is given over about 30 minutes on Day 1 every 3 weeks (21 days). The visit takes longer for assessment and preparation.

STEP 2

Course length

Continue while treatment helps and side effects remain acceptable. At 24 months, the team may consider stopping if there is a sustained response without progression; this is an individual review, not an automatic fixed final cycle.

STEP 3

Blood and thyroid appointments

Blood count, kidney/liver function, glucose and thyroid function at baseline. Blood count, kidney/liver and glucose before each 3-weekly cycle; thyroid every 3–6 weeks regardless of infusion interval; cortisol if indicated. Scans and markers follow the consultant plan.

STEP 4

Home medicines and records

There is no take-home chemotherapy pump. Keep your alert card, symptom record and medicine list. Discuss travel, work and transport needs.

STEP 5

After the final dose

Keep reporting new symptoms. Ask who follows persistent side effects and any hormone medicines.

Sources: NCCP 00455 v16 official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.