Cisplatin + etoposide + chest radiotherapy

Chemotherapy and chest radiotherapy work together to treat lung cancer. Your team will explain the aim, possible benefit and alternatives for you.


Stage III non-small-cell lung cancer · NCCP 00456 v4 · Document revised 7 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

This guide is for Stage III non-small-cell lung cancer. Your radiation team supplies the exact radiation calendar; later treatment is a separate decision.

My overall treatment

Cisplatin on Days 1 and 8; etoposide on Days 1–5. Each cycle lasts 28 days. Two cycles first; two more may be offered after review. Radiotherapy usually starts within 24 hours of the first chemotherapy dose; follow your separate radiation appointment plan.



Chemotherapy + radiotherapy
Two 28-day chemotherapy cycles; radiation runs alongside on its own appointment plan.

Review
Two more cycles only if prescribed after review; not automatic.

After treatment
Recovery checks and the next treatment decision follow your individual plan.

A typical 28-day cycle

Day 1	Day 2–5	Day 6–7	Day 8	Day 9–28
Both chemotherapy medicines	Etoposide each day	No planned chemotherapy	Cisplatin only	No further chemotherapy this cycle

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Blood count, kidneys / liver** Required; extra tests if indicated	Before first treatment; hearing / clearance tests if needed
During	**Blood count** Regimen + etoposide product information	Days 1 and 8; also before each etoposide dose
During	**Kidneys / liver and symptoms** Required safety checks	Before each cycle; extra checks if unwell. Review swallowing and breathing.
After / review	**Recovery, scans and markers** Individual treatment plan	Team-agreed follow-up; no fixed scan or tumour-marker interval in this regimen.

Between treatments

- Tiredness, sickness or swallowing discomfort can occur; report difficulty eating or drinking early.
- Report fever or new breathing changes at any time, including after treatment.

Bring your medicine list and both treatment calendars. Confirm tests, support medicines and urgent contacts before leaving.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hearing / nerves


Cisplatin

Report ringing in the ears, hearing change or new tingling.

2 Mouth / swallowing

Chemotherapy + chest radiotherapy

Report mouth sores, painful swallowing or difficulty drinking.



3 Breathing

Chest radiotherapy + chemotherapy

New or worsening cough or breathlessness needs prompt advice, during or after treatment.

4 Kidneys / fluids

Cisplatin

Report much less urine, swelling or trouble drinking.

5 Blood cells

Chemotherapy

Fever, chills, sudden illness or unusual bleeding need immediate advice.

6 Skin / energy

Chest radiotherapy + chemotherapy

Tell the team about sore skin in the treated area, a spreading rash or increasing fatigue.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call oncology immediately for fever, sudden illness, new or worsening breathlessness, troublesome bleeding, inability to drink or swallow, uncontrolled vomiting, very little urine or severe rash. Tell staff at once about pain, swelling, faintness or breathing trouble during an infusion.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Stage III non-small-cell lung cancer · NCCP 00456 v4 · Document revised 7 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Cisplatin + etoposide + chest radiotherapy (RT)

28-day cycles: Cisplatin D1/8; etoposide D1–5; chest RT. 2 cycles, then review.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

~~EXAMPLE NUMBERS — replace before patient use.~~

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.