Atezolizumab + nab-paclitaxel

Atezolizumab 840 mg is given on Days 1 and 15, before nab-paclitaxel 100 mg/m² on Days 1, 8 and 15 of each 28-day cycle. Nab-paclitaxel is planned for up to six cycles; extension needs consultant review. Atezolizumab may continue after chemotherapy stops.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Breathing

Atezolizumab

New cough or breathlessness may reflect immune inflammation.

3 Liver


Atezolizumab

Jaundice or dark urine needs prompt review.

5 Blood / infection

Nab-paclitaxel

Fever, bruising or sudden illness: contact oncology.



2 Bowel

Atezolizumab / nab-paclitaxel

Report diarrhoea early, especially pain or blood in stool.

4 Hormones / energy

Atezolizumab

Marked weakness, thirst, headache or dizziness needs assessment.

6 Hands / feet

Nab-paclitaxel

Report numbness or tingling before another dose.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects; this is not a complete list. Report new or persistent symptoms even after treatment stops.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Atezolizumab

- Immune inflammation of lungs, bowel, liver or kidneys
- Thyroid, adrenal or pituitary problems and diabetes
- Rare serious heart, nerve or muscle inflammation
- Infusion reactions and severe skin reactions

Nab-paclitaxel

- Numbness / tingling or weakness, sometimes lasting
- Low blood counts, infection and sepsis
- Hair loss, fatigue and muscle / joint aches
- Infusion reactions; uncommon serious lung inflammation

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Know the treatment day

Day 8 is nab-paclitaxel only. Confirm which medicines are due at every visit.

Immune symptoms

Report new symptoms promptly, even after treatment ends. Do not assume diarrhoea or breathlessness is an ordinary chemotherapy effect.

Nerves and blood counts

Report numbness before the next dose; fever or unusual bleeding needs urgent advice.

Medicines and vaccines

Discuss steroids, new medicines and vaccinations with the team.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00688/

Sources and review status are available online. Follow your own prescription and hospital instructions.