# Bevacizumab + weekly paclitaxel

Bevacizumab 10 mg/kg is given on Days 1 and 15, with paclitaxel 80 mg/m² on Days 1, 8 and 15 of each 28-day cycle. Continue according to benefit, toxicity and the treating team's plan.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

## A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

### 1 Blood pressure / vision


**Bevacizumab**

Severe headache or visual change needs urgent review.

### 2 Urine / kidneys

**Bevacizumab**

Protein checks can detect problems before symptoms; report swelling.




### 3 Abdomen / wounds

**Bevacizumab**

Severe abdominal pain or poor wound healing needs urgent advice.

### 4 Bleeding / clots

**Bevacizumab**

Report bleeding; sudden chest, breathing or stroke symptoms are emergencies.

### 5 Blood / infection

**Paclitaxel**

Fever, shivers or sudden illness: contact oncology.

### 6 Hands / feet

**Paclitaxel**

Report tingling or numbness before the next dose.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects; this is not a complete list. Report new or persistent symptoms even after treatment stops.

Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

# Your treatment, at home.

## Important medicine-specific effects

### Bevacizumab

- High blood pressure / urine protein
- Bleeding or blood clots
- Delayed wound healing
- Rare bowel perforation / fistula
- Rare serious neurological effects
- Infusion reaction

### Paclitaxel

- Low blood counts and infection
- Numbness / tingling, sometimes lasting
- Hair loss, muscle or joint pain
- Infusion reaction
- Mouth, bowel, skin and nail effects

**Surgery, radiation and monitoring:** open the whole-treatment map or scan the QR.

## Make the plan practical

### Blood pressure and urine

Attend the required checks even if you feel well. Report severe headache, visual change or swelling.

### Operations and dental work

Tell oncology about planned procedures; bevacizumab timing and wound healing need review.

### Paclitaxel premedicines

The first dose and later doses have different plans. A previous reaction changes the assessment.

### Nerves and bleeding

Report numbness, unusual bleeding or bruising before the next visit.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

**Insert numbers here**

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

**Insert numbers here**

Ask for medical oncology

EVENINGS / OVERNIGHT

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

**Complete reference, visual timeline and passport**

https://sactcheck.com/patient/00738/

Sources and review status are available online. Follow your own prescription and hospital instructions.