# Phesgo + docetaxel

Phesgo and docetaxel on Day 1 every 21 days. Docetaxel is planned for at least six cycles; it may stop after 6–8 cycles or toxicity. Phesgo may continue alone while beneficial and tolerated.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

## A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

### 1 Heart / breathing

**Phesgo**

Keep echoes; report breathlessness, swelling or reduced exercise tolerance.

### 2 Diarrhoea

**Especially pertuzumab**

Report watery stools early, particularly with poor intake or dizziness.

### 3 Injection reaction


**Phesgo**

Tell staff immediately about rash, chest tightness or breathing difficulty.

### 4 Thigh injection site

**Phesgo**

Report persistent or spreading redness, swelling or pain.




### 5 Hands / feet

**Docetaxel**

Report numbness or tingling before the next dose.

### 6 Blood / infection

**Chemotherapy**

Fever, shivers or suddenly unwell: call immediately.

### 7 Hair / scalp

**Chemotherapy**

Hair loss is common; ask about scalp cooling where suitable.

### 8 Mouth / appetite

**Chemotherapy**

Painful ulcers or reduced drinking need advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

The body map follows the medicines active in this phase.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

# Your treatment, at home.

## Important medicine-specific effects

### Docetaxel

- Low blood counts, infection and sepsis
- Hair loss, mouth soreness and fatigue
- Numbness / tingling; sometimes lasting
- Fluid retention and nail changes
- Allergic infusion reaction

### Pertuzumab

- Diarrhoea and dehydration
- Injection / allergic reaction
- Heart pumping-function reduction
- Rash and fatigue

### Trastuzumab

- Heart pumping-function reduction / heart failure
- Injection or allergic reaction
- Diarrhoea and fatigue
- Rare serious lung effects

**Surgery, radiation and monitoring:** open the whole-treatment map or scan the QR.

## Make the plan practical

### Antiemetic plan

Taxane treatment uses a low-risk antiemetic plan; Phesgo alone uses a separate minimal-risk plan.

### Premedicines

Docetaxel: prescribed dexamethasone starts the day before each dose. Ask for one reconciled steroid schedule.

### Appointments and results

Echo before treatment and every 12 weeks (around cycle 5 if on time), including maintenance. Bloods before each cycle. Ask who books and reviews each test; report a missing appointment.

### Phase transition

After chemotherapy stops, use the agreed Phesgo maintenance plan (NCCP 00785 when eligible). Keep cardiac and response checks; chemotherapy steroids, antiemetics and growth factors do not automatically continue.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

**Insert numbers here**

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

**Insert numbers here**

Ask for medical oncology

EVENINGS / OVERNIGHT

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

**Complete reference, visual timeline and passport**

https://sactcheck.com/patient/00796/

Sources and review status are available online. Follow your own prescription and hospital instructions.