Lomustine for recurrent malignant glioma

Treatment to help control recurrent glioma; agree symptom, function and quality-of-life goals with the team.

NCCP 00805: recurrent malignant glioma. · NCCP 00805 v2 · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

NCCP 00805: recurrent malignant glioma. The treating team confirms the applicable indication, suitability and local consent requirements.

My overall treatment

Lomustine once on D1, then no further doses in the 42-day cycle.

Lomustine for recurrent malignant glioma — 42-day view



Lomustine · one oral dose
Day 1

This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

Each cycle reviewed
No fixed total; the team confirms each start and when to stop.

A typical 42-day cycle

Day 1	Day 2–42
One authorised capsule dose	No scheduled lomustine; continue monitoring

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline safety** Regimen / product information	Blood count, kidneys/liver; lung assessment.
During	**Treatment checks** Required by regimen	Regular bloods; review symptoms and prior toxicity before treatment.
Between / after	**Continuing follow-up** Product information / individual plan	Weekly counts ≥6 weeks after lomustine (SmPC); lung follow-up. Individual magnetic resonance imaging (MRI) plan.

Between treatments

- Lomustine is taken once, not daily; counts can fall several weeks later.
- Report fever, bleeding, persistent vomiting, cough or breathlessness promptly.

Bring a clear day-specific medicine plan, anti-sickness/bowel instructions and actual urgent contacts.

When to get help
Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.
Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321
EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Brain / energy


Lomustine / tumour

Fatigue or headache may occur. New seizures, sudden weakness, speech difficulty or loss of consciousness needs emergency help.

2 Mouth / eating

Lomustine

Report mouth soreness or ulcers and reduced intake. Gentle mouth care and early help can support eating and drinking.



3 Blood / infection

Lomustine

Low counts can be delayed for several weeks and build up over cycles. Fever, bruising or bleeding needs urgent advice.

4 Lungs

Lomustine

Lung injury may develop during treatment. Report new cough or breathlessness and attend the lung checks arranged.

5 Liver

Lomustine

Attend liver tests and report yellow skin or eyes, dark urine or unusual itching. Do not assume a rest day means no risk.

6 Kidneys / urine

Lomustine

Kidney checks guide safe treatment. Report reduced urine, swelling or difficulty drinking and follow the team’s fluid advice.

7 Nausea / bowels

Lomustine / support medicines

Take anti-sickness medicine as prescribed. Call for repeated vomiting, inability to drink, constipation or abdominal pain.

8 Fertility / pregnancy

Lomustine

Discuss contraception and fertility before starting. Do not breastfeed during treatment. Follow written product-specific precautions.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Use your oncology unit’s written emergency plan. Fever, shivering or suddenly feeling unwell needs urgent advice even on days without chemotherapy. Call 112 or 999 for a new seizure, sudden weakness or speech difficulty, loss of consciousness, severe breathing difficulty or uncontrolled bleeding. If you have an established seizure rescue plan, follow it and its emergency instructions. Call promptly for fever, bleeding, new cough or breathlessness, persistent vomiting or reduced urine. Do not take another lomustine dose without a new instruction.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital’s temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

NCCP 00805: recurrent malignant glioma. · NCCP 00805 v2 · Document revised 5 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Lomustine · 42 days

Lomustine once on D1, then no further doses in the 42-day cycle.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Ongoing record: the team authorises each cycle; no fixed total is implied.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.