Nivolumab, gemcitabine and carboplatin before surgery

Up to three 21-day cycles. Day 1: nivolumab, gemcitabine and carboplatin. Day 8: gemcitabine alone after blood tests and review. Then surgical multidisciplinary team (MDT) reassessment. The five listed backbones are alternatives; this guide covers only your selected combination.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Hormones / energy

Nivolumab

Report severe tiredness, headache, dizziness or unusual thirst.

2 Breathing

Nivolumab / gemcitabine

Report new cough, breathlessness or chest pain urgently.

3 Liver

Nivolumab / chemotherapy

Blood tests check liver function; report yellow eyes or dark urine.

4 Bowels

Nivolumab / chemotherapy

Report persistent diarrhoea, blood, pain or inability to drink.

5 Kidneys / urine

Nivolumab / carboplatin

Blood tests check kidney function; report reduced urine.

6 Blood / infection


gemcitabine / carboplatin

Fever or sudden illness can be an emergency; follow your contact plan.

7 Skin / hair

gemcitabine / nivolumab

Report a severe rash or blistering urgently; hair changes vary.



Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact your oncology team urgently for fever, feeling suddenly unwell, new breathlessness, chest pain, persistent diarrhoea, severe weakness, confusion, reduced urine or a severe rash. Follow your local emergency instructions; do not wait for the next appointment.

Your treatment, at home.

Important medicine-specific effects

Nivolumab

- Immune inflammation in multiple organs
- Infusion reactions
- Delayed effects after treatment stops

Gemcitabine

- Low blood counts and infection
- Nausea, fatigue and flu-like symptoms
- Skin rash and swelling
- Rare serious lung or kidney injury

Carboplatin

- Low blood counts and infection
- Nausea and vomiting
- Kidney, hearing and nerve effects
- Allergic reactions

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Your written prevention plan

Follow the written anti-sickness plan; Day 8 gemcitabine has its own review and support instructions. Carboplatin dosing uses a verified kidney-function measurement; report dehydration or a change in health.

Immune alert card

Nivolumab can inflame healthy organs during treatment or after it stops. Report new symptoms promptly and carry your immune alert card for at least five months after the last dose.

Prepare for surgery

The oncology and surgical teams agree scans, recovery checks and timing. Report any illness or new symptom before the planned operation.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00849/backbone_00310/

Sources and review status are available online. Follow your own prescription and hospital instructions.