Docetaxel 100 mg/m² · every 3 weeks

Docetaxel 100 mg/m² by intravenous (IV) infusion on Day 1 every 21 days, with dose changes when needed. The number of cycles follows benefit, tolerance and the agreed treatment plan.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Hands / feet

Docetaxel

Report numbness or tingling before the next dose.

3 Hair / scalp


Docetaxel

Hair loss is common. Ask about scalp cooling before treatment where suitable.

5 Mouth

Docetaxel

Painful ulcers or difficulty drinking need advice.



2 Blood / infection

Docetaxel

Fever, shivers or suddenly unwell: call immediately.

4 Fluid / skin / nails

Docetaxel

Report swelling, painful skin or nail changes.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects; this is not a complete list. Report new or persistent symptoms even after treatment stops.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Docetaxel

- Low blood counts, infection and sepsis
- Hair loss, mouth soreness and fatigue
- Numbness / tingling; sometimes lasting
- Fluid retention and nail changes
- Allergic infusion reaction

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Blood counts and infection

Fever, shivers or sudden illness needs prompt advice on any day.

Nerve symptoms

Report numbness, tingling or weakness before the next dose.

Support medicines

Follow the prescription for the active medicine and treatment day.

Review the plan

Discuss symptoms, recovery and the next treatment decision with the team.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00202/

Sources and review status are available online. Follow your own prescription and hospital instructions.