Weekly cetuximab

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Subsequent weekly cetuximab

Active medicines: cetuximab

Emetogenic risk: low

Medicine	Dose / route	When
Chlorphenamine	10 mg intravenous (IV)	60 minutes before each infusion
Dexamethasone	8 mg IV	60 minutes before each infusion Reconcile antiemetic steroid; no automatic duplication.

Additional phase-specific support

- Premedication continues for subsequent infusions.
- Post-treatment monthly electrolytes/Mg/Ca for two months.
- NCCP p1 permits cessation of observation after two tolerated cetuximab doses, whereas Erbitux SmPC section 4.2 requires monitoring during and for at least one hour after each infusion. Reconcile and document the local observation plan; do not promise early discharge.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00207 v6 · p4 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-11-07 · SACTCheck v0.83.8 · Regimen-level drafts; prescriber and oncology-pharmacy verification required