Irinotecan · every three weeks


The aim is to control advanced bowel cancer and symptoms. The team reviews how well treatment is working and whether side effects remain acceptable; there is no single fixed number of cycles for everyone.

Advanced / metastatic colorectal cancer · NCCP 00213 v7 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Irinotecan on Day 1 every 21 days while beneficial and tolerable. There is no home chemotherapy pump. Blood counts are checked weekly, including between infusion visits.



A 21-day reference cycle

Irinotecan on Day 1 every 21 days while beneficial and tolerable. There is no home chemotherapy pump. Blood counts are checked weekly, including between infusion visits.

Your treatment schedule

Day 1	Every week	Through Day 21
Day-ward irinotecan. Report early cramps or sweating immediately.	Full blood count, including between infusions. No home pump.	Use delayed-diarrhoea plan; next treatment only after recovery and review.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline bloods** Required by regimen	Blood count, kidney and liver function before starting.
During	**Weekly full blood count** Required by regimen	Every week, including the two weeks between 21-day infusions.
Before each cycle	**Kidneys / liver / symptoms** Required by regimen	Kidney and liver results, bowel symptoms, hydration and medicines reviewed before more treatment.
After / review	**Recovery / scans / markers** Team-agreed plan	Follow unresolved effects and the individual cancer review plan; no universal scan or marker interval.

Between treatments

- No guaranteed safe days: report fever, sudden illness, diarrhoea or poor intake promptly.
- Symptoms can continue after pump removal and after treatment ends.

Confirm weekly blood-test appointments, anti-sickness medicines, written diarrhoea instructions and working urgent contacts.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hair / scalp

Irinotecan

Hair thinning or loss may occur. Ask about practical support.

2 Mouth / swallowing

Irinotecan

Report ulcers or pain affecting eating or drinking.

3 Breathing / reaction


Irinotecan

New breathlessness, wheeze or severe breathing difficulty needs urgent assessment.

4 Sweating / early reaction

Irinotecan

Cramps, sweating or diarrhoea during infusion: tell the nurse immediately.



5 Bowel / stoma

Irinotecan

Delayed diarrhoea after 24 hours needs your written plan; fever or dehydration needs urgent advice.

6 Hydration / urine

Irinotecan-related fluid loss

Reduced urine, dizziness or inability to replace fluids needs prompt advice.

7 Blood / infection

Irinotecan

Keep weekly blood tests. Fever, shivers or sudden illness: call oncology immediately.

8 Infusion access

Treatment delivery

Report pain, swelling or leakage during infusion. This regimen has no home pump.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

New breathing difficulty, chest pain, collapse, severe diarrhoea or inability to drink require prompt assessment.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Advanced / metastatic colorectal cancer · NCCP 00213 v7 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Irinotecan · every three weeks

Irinotecan Day 1 every 21 days · weekly blood counts · no home chemotherapy pump.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.