# Aflibercept + FOLFIRI

Aflibercept with FOLFIRI is used for metastatic colorectal cancer that is resistant to, or has progressed after, an oxaliplatin-containing treatment. The aim is to control the cancer and symptoms; treatment continues while its benefit outweighs its side effects.

**Metastatic colorectal cancer** · NCCP 00238 v6 · Document revised 4 November 2026

**Your personal plan:** this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

## My overall treatment

Day 1: aflibercept, then irinotecan, folinic acid and fluorouracil, followed by a 46-hour fluorouracil pump. Repeat every 14 days while beneficial and tolerable.




**14-day treatment cycle**

Day 1: aflibercept, then irinotecan, folinic acid and fluorouracil, followed by a 46-hour fluorouracil pump. Repeat every 14 days while beneficial and tolerable.

## Your treatment schedule

| Day 1 | Days 2–3 | Days 4–14 |
|---|---|---|
| Day-ward medicines; pump connected. | Complete 46-hour pump; disconnect as arranged. | Recovery, symptoms and tests; next cycle only after team review. |

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

## My tests and checks

| When | What is checked | Timing / reason |
|---|---|---|
| **Before** | **dihydropyrimidine dehydrogenase (a medicine-processing enzyme) (DPD) / baseline bloods**<br>Required by regimen | DPD before fluoropyrimidine; blood count, kidney and liver tests. |
| **During** | **Bloods**<br>Required by regimen | Blood count, kidney and liver tests before each cycle. |
| **Before / during** | **blood pressure (BP) / urine protein**<br>Required by regimen | Baseline and each-cycle urine protein; BP monitored during treatment; extra urine collection if indicated. |
| **If indicated** | **Heart / international normalised ratio (blood-clotting test) (INR)**<br>Conditional tests | electrocardiogram (ECG) if cardiac function is compromised. If warfarin: INR as clinically indicated. |
| **After / review** | **Recovery / scans / markers**<br>Team-agreed plan | Follow the individual cancer plan and any unresolved effects; scan and marker intervals are team-agreed. |

## Between treatments

- No guaranteed safe days: report fever, sudden illness, diarrhoea or poor intake promptly.
- Symptoms can continue after pump removal and after treatment ends.

Confirm supportive medicines, tests, working urgent contacts and pump disconnection.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**



Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# What to watch for

You do not need to know which medicine caused a symptom before calling.

### 1 Mouth / swallowing

**Fluorouracil + folinic acid**

Report ulcers or pain that affects eating or drinking.

### 3 Bowel / stoma

**Irinotecan / fluorouracil**

Call early for diarrhoea, increased stoma output or poor intake; severe pain needs urgent assessment.

### 5 Blood / infection / bleeding


**Chemotherapy / aflibercept**

Fever, shivers or sudden illness: call now. Major bleeding or a swollen painful limb needs urgent review.

### 7 Head / blood pressure

**Aflibercept**

Severe headache, sudden weakness or visual change needs urgent assessment.




### 2 Chest

**Fluorouracil**

New chest pain, collapse or major breathing difficulty needs emergency help.

### 4 Hands / feet

**Fluorouracil**

Report painful redness, peeling or difficulty with everyday activities.

### 6 Kidneys / urine protein

**Aflibercept**

Keep urine and blood-pressure checks; report swelling or reduced urine.

### 8 Pump / line / healing

**Treatment delivery / aflibercept**

Call for leakage, swelling, line pain or a loose connection. Report poor wound healing and planned procedures.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Fever, shivers or sudden illness needs immediate oncology advice. Chest pain, collapse or severe breathing difficulty needs emergency help. Call urgently for severe or persistent diarrhoea, severe abdominal pain, inability to drink, reduced urine or a line/pump problem. Major bleeding, a painful swollen limb, sudden weakness, a severe headache or visual change also needs urgent assessment.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**


Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# Keep your treatment details to hand

**Metastatic colorectal cancer** · NCCP 00238 v6 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

**FRONT / CONTACTS**

## My Treatment Card

SACTCheck

Name: _____

**Aflibercept + FOLFIRI**

Every 14 days · 46-hour fluorouracil pump · cycle plan confirmed by team.

Phase: _____ Cycle plan: team to confirm

**Daytime (EXAMPLE)** 087 12345678

**Out of hours (EXAMPLE)** 087 87654321

**Unwell or fever? Call oncology now.**
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

**BACK / TREATMENT RECORD**

## My cycle record

Rolling record — follow your agreed treatment plan.

| Cycle | Date / initials |
|---|---|
| | |
| | |
| | |
| | |

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

**FOLD centre — do not cut**

## Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

## Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

### iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

### Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

## Your local contacts and resources

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.



Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.