# Everolimus · neuroendocrine tumours

Daily targeted tablets to help control particular advanced neuroendocrine tumours (NETs).

**Progressive unresectable or metastatic pancreatic neuroendocrine tumour (NET), or well-differentiated G1/G2 nonfunctional gastrointestinal NET under NCCP 00320b/c.** · NCCP 00320 v6 · Document revised 5 November 2026

**Your personal plan:** this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Progressive unresectable or metastatic pancreatic NET, or well-differentiated G1/G2 nonfunctional gastrointestinal NET under NCCP 00320b/c.

## My overall treatment

Once daily every day, consistently with or without food; no planned rest week.

**28** day reference

Reference window · ongoing treatment

**Everolimus every day** — Day 1–28

**Continuous treatment**

Four-week review windows are for monitoring, not a fixed course or routine break.

## A typical 28-day cycle

| Day 1–7 | Day 8–28 |
|---|---|
| Starting safely | Continue daily; report symptoms early |

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

## My tests and checks

| When | What is checked | Timing / reason |
|---|---|---|
| **Before** | **Bloods / urine / infection**<br>Required by regimen | Counts, kidneys/liver, urine protein, glucose, fats and hepatitis B screen. |
| **During** | **Safety bloods**<br>Required by regimen | Counts, kidneys and liver every 4 weeks. |
| **As planned** | **Glucose / fats / NET review**<br>Required / individual timing | Glucose as indicated; fats regularly; scans and hormone markers on your plan. |

## Between treatments

- Report mouth ulcers, new cough, breathlessness, fever or sudden illness promptly.
- Attend tests even if well; do not restart after a hold without instructions.

Swallow whole. Missed dose: no extra tablet; take the next prescribed dose at the usual time.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**



Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# What to watch for

You do not need to know which medicine caused a symptom before calling.

## 1 Mouth / eating


**Everolimus**

Mouth ulcers may appear early. Use prescribed mouth care; report soreness, pain or difficulty eating and drinking.

## 2 Lungs / breathing

**Everolimus**

New cough, fever or breathlessness may be infection or lung inflammation. Contact oncology promptly; severe breathlessness is an emergency.




## 3 Blood / infection

**Everolimus**

Blood counts can fall. Fever, shivering, unusual bruising or sudden illness needs urgent advice even without a scheduled visit.

## 4 Liver

**Everolimus**

Tests can change before symptoms occur. Attend blood checks and report yellow skin or eyes, dark urine or persistent abdominal pain.

## 5 Kidneys / fluid

**Everolimus**

Kidney problems or protein in urine may need extra tests. Report swelling, dehydration or much less urine.

## 6 Bowel / appetite

**Everolimus**

Diarrhoea, nausea or poor appetite can affect fluids and weight. Report persistent symptoms or inability to drink.

## 7 Skin / wounds

**Everolimus**

Rash and slow wound healing can occur. Report severe rash or unhealed wounds and discuss surgery before it is arranged.

## 8 Blood sugar / energy

**Everolimus**

Blood sugar and fats may rise without symptoms. Report marked thirst, frequent urine or unusual fatigue; keep the agreed tests.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact oncology promptly for new or worsening cough, breathlessness or fever, mouth ulcers limiting eating or drinking, marked thirst, much less urine, severe diarrhoea or signs of infection. Severe breathing difficulty or collapse needs emergency help.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**


Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# Keep your treatment details to hand

**Progressive unresectable or metastatic pancreatic NET, or well-differentiated G1/G2 nonfunctional gastrointestinal NET under NCCP 00320b/c.** · NCCP 00320 v6 · Document revised 5 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

**FRONT / CONTACTS**

## My Treatment Card

**SACTCheck**

Name: _____

**Everolimus · neuroendocrine tumours**

Once daily every day, consistently with or without food; no planned rest week.

Phase: _____ Cycle plan: team to confirm

**Daytime (EXAMPLE)** 087 12345678

**Out of hours (EXAMPLE)** 087 87654321

**Unwell or fever? Call oncology now.**
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

**BACK / TREATMENT RECORD**

## My cycle record

Rolling review record; no fixed final cycle.

| Cycle | Date / initials |
|---|---|
| | |
| | |
| | |
| | |

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

**FOLD centre — do not cut**

## Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

## Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

### iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

### Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

## Your local contacts and resources

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE ONLY — team must replace both numbers before patient use.**

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.



Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.