Fortnightly cetuximab + irinotecan

To control advanced bowel cancer and symptoms. Continue while beneficial and tolerable; your team reviews the plan.

Advanced / **metastatic colorectal cancer** · NCCP 00331 v7 · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Cetuximab and irinotecan on Day 1, repeated every 14 days while beneficial and tolerable. There is no Day 8 cetuximab visit and no fluorouracil home pump.

14 day view

Treatment timing in this window

- **Cetuximab infusion** — Day 1
- **Day-ward chemotherapy** — Day 1

A 14-day reference cycle

Cetuximab and irinotecan on Day 1, repeated every 14 days while beneficial and tolerable. There is no Day 8 cetuximab visit and no fluorouracil home pump.

A typical 14-day cycle

Day 1	Day 2–7	Day 8–14
Treatment and checks	Care between visits	Recovery and the next review

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Bloods and suitability** Regimen / treatment suitability	Blood count, kidney/liver tests and tumour RAS suitability. Review medicines, allergies and symptoms.
During	**Bloods, skin and symptoms** Required by regimen	Regular blood count, kidney/liver tests as arranged; check skin, eyes, bowel symptoms and hydration before treatment.
During / after	**Magnesium and other minerals** Required by regimen	Monitor during treatment; monthly electrolytes, magnesium and calcium for two months after the last antibody dose.
Review	**Response and recovery** Individual plan	Scans and other response tests follow the consultant plan; follow unresolved effects.

Between treatments

- Report cramps or sweating during infusion. At the first liquid stool, use your supplied rescue plan, drink advised electrolyte fluids and contact the team early. If no plan was supplied, call immediately.
- Fever, severe diarrhoea, poor intake or reduced urine needs urgent advice. Symptoms can occur on any day.

Bring your medicine list and report new symptoms before dosing. Confirm test appointments, the skin-care plan and urgent contacts. Have a written irinotecan diarrhoea rescue plan. Report symptoms before any restart; never adjust doses yourself.

When to get help

Call 112 / **999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Eyes

Cetuximab

Eye pain or new visual change needs prompt advice.

2 Mouth / swallowing

Treatment

Pain or ulcers affecting eating or drinking need advice.

3 Breathing / reaction


Cetuximab

New breathing difficulty requires prompt assessment.

4 Chest

Treatment

New chest pain needs urgent assessment.



5 Bowel

Irinotecan

Diarrhoea, increased stoma output or dehydration: call early.

6 Skin / nails

Cetuximab

Report painful rash, cracks or infected skin early.

7 Blood / infection

Treatment

Fever, shivers or sudden illness: contact oncology immediately.

8 Infusion access

Treatment delivery

Pain, swelling, leakage or a loose connection needs immediate advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

New breathing difficulty, chest pain, collapse, severe diarrhoea or inability to drink require prompt assessment. Report eye pain, blurred vision, blistering or infected skin promptly.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Keep your treatment details to hand

Advanced / metastatic colorectal cancer · NCCP 00331 v7 · Document revised 5 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Fortnightly cetuximab + irinotecan

Cetuximab + irinotecan D1 every 14 days; no D8 visit or home pump.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record; the team confirms the total treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.