Nintedanib + docetaxel


To help control adenocarcinoma lung cancer after first-line chemotherapy; treatment benefit and side effects are reviewed together.

Adenocarcinoma non-small-cell lung cancer (NSCLC) · **combination phase** · NCCP 00372 v6 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Day 1: docetaxel drip, with no nintedanib. Days 2–21: nintedanib capsules twice daily. Repeat every 21 days for up to 6 cycles (about 18 weeks without delays); treatment may stop earlier. A later tablet-only phase needs a new team-confirmed plan.



Combination phase
Day 1 docetaxel; Days 2–21 nintedanib twice daily. Up to 6 cycles.

Possible tablet-only phase
Only after at least 4 completed combination cycles and team approval: nintedanib every day in 28-day cycles. This change is not automatic.

A typical 21-day cycle

Day 1	Day 2–21
Docetaxel day	Nintedanib at home

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Blood count, kidney/liver tests and blood pressure (BP)** Required by regimen	Before treatment; confirm allergy, medicines and pregnancy-prevention advice
During	**Blood count, kidney/liver tests** Required by regimen	Before each 21-day cycle; blood count also around the expected low-count period
During	**Symptoms and blood pressure** Required by regimen	BP periodically/as needed; extra blood or liver tests for symptoms or risk
After / review	**Phase, blood counts and scans** Source blood-count check; other plan team-agreed	After the last combination cycle: blood count as indicated; continuation requires review; scans/markers individually planned

Between treatments

- Diarrhoea, nausea and fatigue may occur at home. Call early if drinking becomes difficult or symptoms persist.
- Low blood counts can occur between visits. The team will set an extra blood-test date; there is no guaranteed safe day.

Bring your medicine list and report new symptoms before dosing. Confirm test appointments, the current phase and daytime / out-of-hours contact plan.

When to get help
Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.
Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321
EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Nerves / neurological symptoms

Both medicines

Report tingling; new severe headache, confusion, seizure or visual change is urgent.

3 Blood vessels / blood pressure


Nintedanib

Chest pain, sudden breathlessness, bleeding or a painful swollen leg needs urgent help.

5 Bowels / sickness

Both medicines

Call early for diarrhoea or vomiting; severe pain needs urgent assessment.



2 Mouth

Docetaxel

Report ulcers or pain that limits eating or drinking.

4 Liver

Nintedanib and docetaxel

Attend liver tests; report yellow skin/eyes or dark urine.

6 Blood counts / infection

Docetaxel with nintedanib

Fever, shaking chills or sudden illness needs immediate oncology advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call your oncology team immediately for fever, shaking chills, sudden illness, breathing difficulty, chest pain, significant bleeding, severe tummy pain, uncontrolled vomiting or new severe headache, confusion, seizure or vision change. Do not wait for the next appointment.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Adenocarcinoma NSCLC · combination phase · NCCP 00372 v6 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Nintedanib + docetaxel

21-day cycle: docetaxel Day 1; nintedanib twice daily Days 2–21. Up to 6 cycles.

Phase: _____ Planned cycles: 6

Daytime (EXAMPLE) 087 12345678
Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Team stamps after each completed cycle.

C1	C2	C3
Date / initials	Date / initials	Date / initials
C4	C5	C6
Date / initials	Date / initials	Date / initials

My treatment information

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.