Panitumumab + FOLFIRI

To control advanced bowel cancer and symptoms. Your team reviews benefit, side effects and the ongoing plan.

RAS wild-type metastatic colorectal cancer; first-line or selected second-line · NCCP 00448 v5 · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Panitumumab and FOLFIRI on Day 1 every 14 days; a 46-hour fluorouracil pump ends around Day 3. No Day 8 infusion.



A repeating 14-day cycle

FOLFIRI includes irinotecan, folinic acid and fluorouracil.

A typical 14-day cycle

Day 1	Day 2–3	Day 4–7	Day 8–14
Infusion and pump connection	The pump is delivering treatment	After disconnection	Between visits and the next review

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Suitability and bloods** Required by regimen	Confirm RAS result, blood count, kidney/liver tests; dihydropyrimidine dehydrogenase (a medicine-processing enzyme) (DPD) before fluorouracil unless already tested.
During	**Bloods and symptom review** Required by regimen	Blood count, kidney/liver tests before each cycle; weekly counts if delayed for low counts. Review skin, bowel and hydration symptoms.
During / after	**Mineral levels** Required by regimen	Monitor during treatment; monthly electrolytes, magnesium and calcium for two months after the last antibody dose.
If indicated	**electrocardiogram (ECG) / warfarin international normalised ratio (blood-clotting test) (INR)** Conditional tests	ECG if cardiac function is compromised; INR as clinically indicated if taking warfarin.
Review	**Response and recovery** Individual plan	Scans and markers follow the consultant plan. Follow unresolved effects.

Between treatments

- Report cramps or sweating during infusion. At the first liquid stool, use your prescribed rescue plan, drink advised electrolyte fluids and contact the team early.
- Fever, sudden illness, severe diarrhoea, poor intake or reduced urine needs urgent advice. There are no guaranteed safe days.

Bring your medicine list; confirm tests, the skin-care plan and working urgent contact numbers. Confirm your pump-care instructions, professional disconnection and written symptom rescue plan.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Eyes

Panitumumab

New eye pain or visual change needs prompt advice.

2 Mouth

Fluorouracil + folinic acid

Report ulcers or pain affecting eating and drinking.

3 Chest

Fluorouracil

New chest pain or breathlessness needs urgent assessment.

4 Bowel

Irinotecan / fluorouracil

Report diarrhoea, increased stoma output or severe abdominal pain.

5 Hydration / minerals

Panitumumab + chemotherapy

Reduced urine, cramps or weakness need advice; keep your blood-test appointments.

6 Skin / nails

Panitumumab

Report rash, cracks or painful nails early.

7 Hands / feet


Fluorouracil

Report painful redness, peeling or difficulty using your hands or walking.

8 Pump / line

Treatment delivery

Pain, leakage, swelling or a loose connection needs immediate advice.



Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Report new breathing difficulty, eye pain or blurred vision, blistering or infected skin, severe diarrhoea or dehydration promptly. Also report new chest pain, fever or a line/pump problem immediately.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

RAS wild-type metastatic colorectal cancer; first-line or selected second-line · NCCP 00448 v5 · Document revised 5 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Panitumumab + FOLFIRI

D1 every 14 days; 46-hour fluorouracil pump to D3; no D8 visit.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record; the team confirms the total treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.