Durvalumab 1500 mg every 28 days

Immunotherapy after chemotherapy and radiotherapy, aiming to reduce the chance of lung cancer returning or worsening.

Stage III non-small-cell lung cancer (NSCLC) · **consolidation after concurrent chemoradiotherapy** · NCCP 00655 v3a · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Durvalumab is given on Day 1 every 28 days, for up to 12 months (13 cycles). Treatment may stop earlier after review.

28 day view

Treatment timing in this window

Durvalumab · Day 1

Day 1

Before durvalumab	Start and continue	After the course
Concurrent chemotherapy and radiotherapy are completed first.	First dose within 42 days after chemoradiotherapy; then every four weeks.	Keep reporting symptoms and follow your monitoring plan.

Your 28-day treatment cycle

Day 1	Days 2–28
Durvalumab infusion after review.	Symptom monitoring and agreed tests; no durvalumab tablets.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Blood count, kidneys, liver, glucose, thyroid, hepatitis B/C and PD-L1** Required by regimen	Baseline checks and review of recovery after chemoradiotherapy
During	**Blood count, kidneys, liver, glucose and thyroid** Required by regimen	Regular testing; your team confirms the dates
After / review	**Recovery and cancer assessment** Team-agreed plan	Timing agreed with your team; report new symptoms promptly

Between treatments

- No durvalumab tablets are taken between infusions.
- Report new cough, breathlessness, diarrhoea, rash or unusual weakness; immune effects can occur after treatment.

Keep your immunotherapy alert card and contact details. Confirm the blood-test and infusion dates.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Lungs / breathing

Durvalumab

Report new or worsening cough, breathlessness or chest pain promptly, including after finishing radiotherapy.

2 Hormones / head / nerves

Durvalumab

Severe headache, dizziness, confusion, unusual weakness or thirst need prompt assessment; collapse or seizures need emergency

3 Heart / muscles

Durvalumab

Chest pain, palpitations, marked weakness or trouble swallowing can signal serious inflammation; seek urgent help.

4 Liver

Durvalumab

Report yellow eyes/skin, dark urine or significant upper abdominal pain; blood tests may detect changes before symptoms.

5 Bowel

Durvalumab

Report new diarrhoea, blood in stools or significant abdominal pain; do not assume this is ordinary stomach upset.

6 Kidneys / urine


Durvalumab

Report much less urine, swelling or a change in urine; attend kidney blood tests.

7 Skin

Durvalumab

Report a spreading rash. Blistering, peeling or mouth/eye sores need urgent assessment.



Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact the oncology team promptly for new or worsening cough, breathlessness or chest pain; diarrhoea, blood in stools or significant abdominal pain; yellow skin/eyes or dark urine; reduced urine; a spreading or blistering rash; unusual weakness, severe headache, dizziness, marked thirst or passing much more urine. Severe breathing difficulty, collapse, confusion, seizures, rapidly worsening weakness or difficulty swallowing need emergency assessment. Do not wait for the next infusion.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Stage III NSCLC · consolidation after concurrent chemoradiotherapy · NCCP 00655 v3a · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Durvalumab

Day 1 every 4 weeks, after chemoradiotherapy. Up to 12 months / 13 cycles.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.