Adult treatment scope · National Cancer Control Programme (NCCP) 00702a / 00702b This adult suite covers entrectinib alone for ROS1-positive advanced non-small-cell lung cancer (NSCLC) before any ROS1 inhibitor (00702a), or a separate NTRK gene-fusion solid-tumour indication with no previous NTRK inhibitor and no satisfactory treatment options (00702b). Molecular tests and Irish access criteria must be confirmed. The adult 600 mg schedule does not apply to children.

NCCP 00702 · VERSION 2


Entrectinib monotherapy · adult

The usual adult starting dose is 600 mg by mouth once daily, every day without a planned rest week. Use your prescribed dose. Treatment continues while helping and side effects remain acceptable; there is no fixed number of cycles.

Oral targeted therapy · selected medicine-specific effects. Full details and sources are in the complete reference.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.



1 Dizziness / thinking changes

entrectinib

Report new confusion, memory or balance problems; avoid driving while affected.

2 Visual changes

entrectinib

New blurred vision or other visual disturbance needs review.

3 Breathlessness / swelling / fainting

entrectinib

Heart pumping or rhythm problems require urgent assessment.

4 Jaundice / liver tests

entrectinib

Liver injury may need blood tests and interruption.

5 Bowel symptoms / nausea

entrectinib

Report severe symptoms or dehydration promptly.

6 Bone pain / altered gait

entrectinib

Possible fracture needs prompt assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call the oncology team urgently for new or worsening breathlessness, swelling, palpitations or fainting; marked confusion or memory change; fever, jaundice, severe diarrhoea or inability to drink. Report new visual disturbance and focal bone pain promptly. Follow your unit's temperature/contact instructions. Severe breathing difficulty, chest pain or collapse needs emergency help. Severe chest pain, major breathing difficulty, collapse or stroke symptoms: 112 / 999.

YOUR PRACTICAL TREATMENT PLAN

Know the next step.

Taking treatment

The usual adult starting dose is 600 mg by mouth once daily, every day without a planned rest week. Use your prescribed dose. Treatment continues while helping and side effects remain acceptable; there is no fixed number of cycles.

Take whole capsules with or without food; do not crush or chew. Avoid grapefruit, grapefruit products and Seville oranges. If a dose is missed, take it unless the next dose is due within 12 hours; confirm the exact boundary with your pharmacist. The leaflet permits repeating whole capsules after immediate vomiting: follow your written plan and call if uncertain. Suspension or tube administration requires separate pharmacy instructions.

Checks and follow-up

Blood count, kidney/liver tests, uric acid and electrocardiogram (ECG)/electrolytes before starting. Repeat ECG/electrolytes after one month, then periodically as indicated; regular blood tests and symptom reviews. Heart-pumping tests depend on symptoms/risk. Scans follow your Consultant's plan. Confirm who arranges and reviews each check.

The whole-treatment map shows the checks for this exact treatment setting.

Support between reviews

Symptom medicines

NCCP rates entrectinib minimal-to-low oral emetic risk. Agree individual rescue treatment and check heart-rhythm interactions; no routine infusion premedication is specified.

Daily life and medicines

Keep the medicine list current. Avoid driving or machinery while dizzy, faint, visually affected or cognitively affected. Review falls risk and report bone pain.

Pregnancy and breastfeeding

Use the agreed contraception plan during and after treatment. Do not breastfeed during treatment; ask about fertility and family plans before starting.

Oncology daytime: Insert numbers here. **Out of hours:** Insert numbers here. Follow your own hospital's urgent-care plan. Emergency: 112 / 999.

Your complete treatment reference

https://sactcheck.com/patient/00702/

Public information only. No personal information is encoded.